

NOTES ON GEOGRAPHIC DISTRIBUTION

Reptilia, Squamata, Teiidae, *Teius suquiensis*: New evidence of recent expansion of this parthenogenetic lizard?

Mario R. Cabrera¹
Julio C. Monguillot²

¹ CONICET and Departamento de Diversidad Biológica y Ecología, FCEFYN, Universidad Nacional de Córdoba Vélez Sarsfield 299, (5000) Córdoba, Argentina. E-mail: mcabrera@efn.uncor.edu

² Administración de Parques Nacionales - Delegación Regional Centro Avenida Richieri 2298, (5000) Córdoba, Argentina.

Teius suquiensis is the only parthenogenetic teiid lizard known in southern South America. From its discovery in central Córdoba province, Argentina, and description (Avila and Martori 1991), new findings, both in museum collections and in the field, have extended its range to the neighboring provinces of San Luis in the southwest (Martori and Aun 1995) and Santa Fe, in the east (Avila 1995). In many localities, the species lives in sympatry with one of the bisexual species, either *Teius teyou* or *T. oculatus*. Sympatric occurrence of the three species has been reported at Daniel Donovan and San Luis city, both in San Luis province (Guerreiro et al. 1998), and between the localities of Gregoria Perez de Denis and Tostado, in Santa Fe province (Avila 2002).

The significant environmental alterations introduced by human activities, particularly deforestation and woodland substitution by soybean and other cultures, have had a profound effect on the lizards, causing the loss of some populations and the possible range expansions of others. *Teius suquiensis*, as well as *T. oculatus*, is frequently associated with disturbed, ecotonal or marginal habitats (Avila and Martori 1991). Also the possibility of ongoing hybridization between the bisexual species of *Teius* has been suggested (Penafort et al. 1986; Avila 2002). In such a dynamic system, it is important to collect any new information able to improve the knowledge on the current ranges of the respective species.

In this paper we report a new case of sympatry between *Teius suquiensis* and *T. teyou*, and extend the range of the former.

All specimens were taken in the surroundings of the ranch El Cercado, 12 km SE of Pozo Nuevo, Department of Sobremonte, Province of Córdoba, Argentina (29°38'39" S, 64°00'51.6" W), on 25 January 2007, between 12:00 and 14:30 h. The collection site is situated on the Sierra de Ambargasta, at 597 m of altitude. Phytogeographically it belongs to the Chaco Formation (*provincia fitogeográfica chaqueña*), within a landscape of quebracho blanco (*Aspidosperma quebracho-blanco*) and *Prosopis* spp. trees, degraded and altered by wood extraction, fires, and cattle raising, that have produced its present physiognomy of secondary thorn woodland (*matorrales de sustitución*) with prevalence of bushes of the genera *Mimozyanthus*, *Celtis*, *Mimosa*, and *Acacia* (Prado 1993; Cabido and Zak 1999).

Three voucher specimens of *Teius suquiensis* (Figure 1) were collected and housed at the herpetological collection of the Comparative Anatomy Laboratory, Universidad Nacional de Córdoba, Argentina, under the catalogue numbers C.484, C.486, and C.488. The only other lizard syntopic with *T. suquiensis* was *T. teyou* (Figure 2), of which we collected two adult females (C.483 and C.487) and one subadult male (C.485).

NOTES ON GEOGRAPHIC DISTRIBUTION



Figure 1. Adult of the all-female species *Teius suquiensis* from Ranch El Cercado, Department of Sobremonte, Córdoba, Argentina.



Figure 2. Adult female of *Teius teyou*, syntopic and sympatric with *T. suquiensis* at Ranch El Cercado.

NOTES ON GEOGRAPHIC DISTRIBUTION




Figure 3. Geographic distribution of *Teius suquiensis*. White circles are data from literature (Avila 2002); black circle represents El Cercado record.

NOTES ON GEOGRAPHIC DISTRIBUTION

The lizards were active on both man-made clearings and sandy secondary roads. Upon pursuit, they rapidly fled, seeking shelter in nearby *Lagostomus* burrows or under bushes. If captured, they vigorously struggled and tried to bite. Body temperature was not measured but the lizards were perceptibly warm. In two complete days of field work only one subadult male (snout-vent length = 105 mm, still without blue pattern) of *Teius teyou* was seen. All others were either adult females of both species or juveniles probably in their second or third year of age, according to their body sizes. At dissection, the females of both species showed big intrabdominal fat bodies, and small ovarian follicles. This agrees with the results of Martori and Aun (1993), who found enlargement of fat bodies after the end of the reproductive season in late December, in a population of *Teius suquiensis* from Bialeto Massé, 190 km to the south from El Cercado.

Other lizards we found occupying different habitats, but not far away, were the common Chacoan iguanids *Tropidurus etheridgei* and *T. spinulosus*, on big rocks, trees, and roof and walls, both inside and outside of the house, and *Stenocercus doellojuradoi* under low bushes with sandy soil. Also an adult Red Tegu lizard (*Tupinambis rufescens*) was seen foraging on the sandy road, a hundred meters away from the house.

The enhanced reproductive potential of an all-female population of lizards as *Teius suquiensis* could provoke the rapid replacement or displacement of sexual species by this partheno-

genetic form. In samples collected by us 23 years ago at the same locality, the only species of *Teius* present was *T. teyou*. Avila (1995) found sympatry of the three species at Santa Fe, in museum samples collected more than 25 years before at the locality of Tostado. However, in a 1995 field trip, he found only *T. suquiensis* at this locality.

The present new record for *Teius suquiensis* extends the previously known range in the province of Córdoba ca. 100 km to the north, into the Chaco Formation, and 220 km to the WSW from the nearest eastern locality, Tostado (Avila 2002). This range extension still does not fill the considerable gap between the populations of Córdoba and Santa Fe provinces (Figure 3). It is important to note that the species has been found neither in an intervening zone with Chaco remnants in northeastern Córdoba (Briguera et al. 2006) nor in the province of Santiago del Estero. Being *T. suquiensis* not typical in Chacoan lizard communities, future work in both unaltered and disturbed Chaco sites would help to better understand the species distribution trend.

Acknowledgements

We are grateful to Norman Scott Jr. and Cristiano Nogueira for comments that improved the manuscript, and Bernardo Sonzini for field companion and logistics. To Lucía Ruiz and Jael Dominino, from the Administración de Parques Nacionales, for their contributions. Financial support was provided by CONICET and SECyT-UNC.

Literature cited

- Avila, L. J. 1995. *Teius suquiensis*: Ampliación de su distribución geográfica y primera cita para la provincia de Santa Fe. Cuadernos de Herpetología 9(1): 57-58.
- Avila, L. J. 2002. Geographic distribution of lizards of the genus *Teius* (Squamata: Teiidae: Teiinae) in southern South America. Biogeographica 78(1): 15-33.
- Avila, L. J. and R. A. Martori. 1991. A unisexual species of *Teius* Merrem 1820 (Sauria Teiidae) from central Argentina. Tropical Zoology 4: 193-201.
- Briguera, V., D. Tamburini, M. Kufner, G. Gavier, L. Giraudo, R. Torres, and V. Bechara. 2006. Herpetofauna en relictos de bosque chaqueño de la región de Mar Chiquita, Córdoba. Cuadernos de Herpetología 20(1): 25-31.
- Cabido, M. R. and M. R. Zak. 1999. Vegetación del Norte de Córdoba. Privately printed, Córdoba. 56 p.
- Guerreiro, A. C., J. C. Baldoni, A. M. Brigada, and M. R. Cabrera. 1998. Ampliación de la distribución de *Teius oculatus* y *T. suquiensis* (Sauria: Teiidae) en la provincia de San Luis (República Argentina). Cuadernos de Herpetología 12(1): 51.
- Martori, R. A. and L. Aun. 1993. Reproductive cycle of the parthenogenetic lizard *Teius suquiensis*. Amphibia-Reptilia 14: 389-393.

NOTES ON GEOGRAPHIC DISTRIBUTION

Martori, R. and L. Aun. 1995. Relevamiento de la herpetofauna de las Sierras Pampeanas del este en la provincia de Córdoba (Argentina) (registro de taxa y comentarios taxonómicos). *Acta Zoologica Lilloana* 43(1): 87-97.

Penafort, M. B., J. A. Scolari, J. M. Cej, and S. Claver. 1986 ("1983"). La identidad de las especies de *Teius* de Argentina y reconocimiento de híbridos en áreas antropizadas de San Rafael, Mendoza, por medio del

análisis discriminante. *Revista del Museo de Historia Natural de San Rafael* 8(4): 109-120.

Prado, D. E. 1993. What is the Gran Chaco vegetation in South America? I. A review. Contribution to the study of flora and vegetation of the Chaco. V. *Candollea* 48(1): 145-172.

Received February 2007

Accepted May 2007

Published online July 2007