

NOTES ON GEOGRAPHIC DISTRIBUTION

Amphibia, Anura, Amphignathodontidae, *Gastrotheca albolineata*: Distribution extension, new state, and new altitudinal records.

Eliana Gressler¹
Andrea G. Aguirre¹
Célio F. B. Haddad²

¹Departamento de Botânica, Instituto de Biociências, Universidade Estadual Paulista.
Caixa Postal 199. CEP 13506-970. Rio Claro, SP, Brazil.

²Departamento de Zoologia, Instituto de Biociências, Universidade Estadual Paulista.
Caixa Postal 199. CEP 13506-970. Rio Claro, SP, Brazil. E-mail: haddad@rc.unesp.br

Gastrotheca albolineata is a medium-size amphignathodontid treefrog described as *Hyla albolineata* by Lutz and Lutz (1939). Sachsse et al. (1999) included this species in the genus *Gastrotheca* based on morphology and reproductive mode, providing a new diagnosis and notes on natural history. The present known distribution of this species is limited to 200–1,400 m above sea level in the states of Rio de Janeiro and Espírito Santo, Brazil (Sachsse et al. 1999; Caramaschi and Rodrigues 2007; Frost 2007). The occurrence of the species in the state of Espírito Santo (Frost 2007) may be incorrect and in the last review for the species of *Gastrotheca* from the Atlantic forest, *G. albolineata* is considered restricted to Serra dos Órgãos, in the state of Rio de Janeiro (Caramaschi and Rodrigues 2007).

During field work at Parque Estadual da Serra do Mar, Núcleo Picinguaba (ca. 23°22' S, 44°48' W; 90 m a.s.l.), municipality of Ubatuba, state of São Paulo, southeastern Brazil, on 27 March 2007, one adult of *Gastrotheca albolineata* was photographed (Figure 1). The vegetation at Núcleo Picinguaba is composed by coastline, dune vegetation, mangroves, coastal plain forest, lower montane or premontane forest (Atlantic rain forest), and montane or cloud forest (SMA 1992). The specimen of *G. albolineata* was photographed

on the forest ground, in a hillside near (200 m) to Rio da Fazenda, ca. 4.0 km from the sea, in the premontane forest (Oliveira-Filho and Fontes 2000). In Picinguaba this type of vegetation presents an irregular canopy (trees 15–30 m high), with high diversity of tree and understudy species; the most-rich families are Myrtaceae, Fabaceae, Rubiaceae and Lauraceae (see detailed description in Sanchez et al. 1999; and Morellato et al. 2000). This first record of *G. albolineata* to the state of São Paulo extends the distribution of the species in approximately 190 km southwest from its previous southern most record (at Petrópolis, state of Rio de Janeiro), and is the lowest altitudinal record for this species. The observed specimen was not collected, being the photograph enough for the proper identification.

Acknowledgments

Ivan Sazima helped with the identification of *G. albolineata*. Renato S. Bérnils made useful comments on the distribution of the species. We also thank Instituto Florestal for work facilities and field lodging at Núcleo Picinguaba, Parque Estadual da Serra do Mar. Eliana Gressler received a fellowship from FAPESP (proc. 05/54267-1) and Célio F. B. Haddad is grateful to FAPESP and CNPq for financial support.

NOTES ON GEOGRAPHIC DISTRIBUTION



Figure 1. Adult of *Gastrotheca albolineata* photographed at Picinguaba, municipality of Ubatuba, state of São Paulo, Brazil

Literature cited

- Caramaschi, U. and M. T. Rodrigues. 2007. Taxonomic status of the species of *Gastrotheca* Fitzinger, 1843 (Amphibia, Anura, Amphignathodontidae) of the Atlantic rain forest of eastern Brazil, with description of a new species. *Boletim do Museu Nacional*, n.s., 525: 1-19.
- Frost, D. R. 2007. Amphibian Species of the World: an Online Reference. Version 5.1 (10 October, 2007). Electronic database accessible at <http://research.amnh.org/herpetology/amphibia/index.php>. American Museum of Natural History, New York, USA. Captured on January 2008.
- Lutz, A. and B. Lutz. 1939. New Hylidae from Brazil. *Anais da Academia Brasileira de Ciências* 11: 67-89.
- Morellato, L. P. C., D. C. Talora, A. Takahasi, C. C. Bencke, E. C. Romera, and V. B. Zipparro. 2000. Phenology of Atlantic rain forest trees – a comparative study. *Biotropica* 32(4B): 811-823.
- Oliveira-Filho, A. T. and M. A. Fontes. 2000. Patterns of floristic differentiation among Atlantic forests in southeastern Brazil, and the influence of climate. *Biotropica* 32: 793-810.
- Sachsse, R., E. Izecksohn, and S. P. de Carvalho-e-Silva. 1999. The systematic status of *Hyla albolineata* Lutz and Lutz, 1939 (Amphibia: Anura: Hylidae). *Herpetologica* 55: 401-406.
- Sanchez, M., F. Pedroni, H. F. Leitão-Filho, and O. César. 1999. Composição florística de um trecho de floresta ripária na Mata Atlântica em Picinguaba, Ubatuba, SP. *Revista Brasileira de Botânica* 22(1): 31-42.
- SMA (Secretaria do Meio Ambiente) 1992. Serra do Mar: uma viagem a Mata Atlântica. São Paulo, Brasil. Secretaria do Meio Ambiente. 94 p.

Received November 2007

Accepted January 2008

Published online January 2008