

First record of the common house gecko *Hemidactylus frenatus* Schlegel, 1836 and distribution extension of *Phyllodactylus reissii* Peters, 1862 in the Galápagos

Omar Torres-Carvajal^{1,*} and Washington Tapia²

1 Pontificia Universidad Católica del Ecuador, Escuela de Biología. Avenida 12 de Octubre y Roca, Apartado 17-01-2184. Quito, Ecuador.

2 Parque Nacional Galápagos. Puerto Ayora, Galápagos, Ecuador

* Corresponding author. E-mail: omartorcar@gmail.com

ABSTRACT: Among introduced species in the Galápagos are three species of geckos – *Gonatodes caudiscutatus*, *Lepidodactylus lugubris*, *Phyllodactylus reissii* – occurring on the islands of Isabela, San Cristóbal, and Santa Cruz. Here we report the first record of a fourth invasive species of gecko from Isabela, as well as the first record of *P. reissii* from the same island.

Invasive species represent one of the major threats to the unique biodiversity of the Galápagos. Some invasive species are conspicuous and obviously harmful (*e.g.*, goats, blackberries), and many efforts have been made to eradicate them (Tapia *et al.* 2000; Carrión *et al.* 2008). Others, however, are small or cryptic (*e.g.*, cockroaches, fire ants), and often difficult to detect (Lubin 1984; Peck *et al.* 1998). The impact caused by these small invaders represents a major challenge, and therefore, prevention or eradication programs are more difficult to establish. Among the small invaders are three species of geckos (Bungartz *et al.* 2011), the only introduced reptiles with established populations in the Galápagos, which share the islands with six endemic species of geckos (Table 1). More than a century ago, the first record of an introduced gecko – *Gonatodes caudiscutatus* Günther 1859 – from San Cristóbal Island was published, although it was originally described as a different species – *G. collaris* – by Garman (1892; synonymy *vide* Vanzolini 1965). This species is widely distributed along the Pacific versant of the Andes in Colombia, Ecuador, and Peru (Burt 1932; Torres-Carvajal 2001; P. Venegas personal communication). While working on the reptiles of the Galápagos Garman received a small collection from Guayaquil (mainland Ecuador), and according to Van Denburgh (1912) it remains unclear whether the specimens of *G. caudiscutatus* examined by Garman were collected in the Galápagos or Guayaquil. Nonetheless, the occurrence of this species in San Cristóbal has been subsequently confirmed (*e.g.*, Mertens 1963; Wright 1983; Olmedo and Cayot 1994). Another introduced species is *Phyllodactylus reissii* Peters 1862, which occurs naturally along the Pacific coast of southern Ecuador and northwestern Peru (Dixon and Huey 1970). This species was found around 1975 in Puerto Ayora, Santa Cruz Island, and was confirmed as a well-established species in that town in 1988 (Hoogmoed 1989). The third known species of an introduced gecko is *Lepidodactylus lugubris* Duméril and Bibron 1836, a parthenogenic species from

the Western Pacific that has invaded many Pacific islands and the Pacific coast of Central and South America. In the Galápagos, this species has been introduced to Santa Cruz, San Cristóbal, and Isabela (Hoogmoed 1989; Olmedo and Cayot 1994). Herein we report the first records of a fourth species of invasive gecko new to the Galápagos, as well as a new island record of *P. reissii*.

During a field collecting trip to Puerto Villamil (0°57'26.42" S, 90°58'2.56" W, WGS84, 0 m), Isabela Island, between 6–8 January 2011, three individuals of *Hemidactylus frenatus* (QCAZ 11128, 11164–65) and two of *Phyllodactylus reissii* (QCAZ 11186–7) were hand captured. All specimens were collected under collection permit PC-05-10 issued by Galápagos National Park, and were deposited at Museo de Zoología QCAZ of Pontificia Universidad Católica del Ecuador (PUCE). Specimens of both species (Figure 1) were found active between 20:00 h–23:00 h on house walls at 1–3 m above ground, usually in areas exposed to artificial light. Although two more specimens of each species were sighted, they were largely outnumbered by *Lepidodactylus lugubris* (Figure 1), which seemed to prefer light posts and palm trees. On a single night (January 6th), 30 specimens of *L. lugubris* were captured by five people between 21:00 h–23:00 h in a two block area of Puerto Villamil. During the same collecting period 24 specimens of the endemic species *P. galapagoensis* were found (Figure 1); they seemed to prefer darker areas closer to the ground, such as lava rock piles.


Without further studies monitoring the species of introduced geckoes on Isabela, it remains unclear whether the specimens of *Hemidactylus frenatus* and *Phyllodactylus reissii* reported here were part of well-established populations. The common house gecko *H. frenatus* has its natural range in tropical Asia and the Indo-Pacific (Case *et al.* 1994), and has been introduced in many tropical and subtropical regions worldwide (Bauer and Henle 1994) including the Pacific coast of Ecuador (Jadin *et al.* 2009),

were it seems to be expanding at a rapid rate (Carvajal-Campos and Torres-Carvajal 2010). Even though nothing is known about the impacts of introduced *H. frenatus* on other species, its seemingly good dispersal ability represents a potential threat for endemic species in Isabela and other islands. The presence of *P. reissii* in Isabela is not surprising, as this species is well-established in Santa Cruz


and there is intensive human and cargo traffic between both islands. The last time that the gecko fauna of Puerto Villamil was thoroughly sampled was in the early 1990s, and *L. lugubris* was the only introduced species reported (Olmedo and Cayot 1994). Therefore, both *H. frenatus* and *P. reissii* have probably arrived on Isabela sometime during the last two decades.

TABLE 1. Origin, activity and distribution (Van Denburgh 1912; Lanza 1973; Olmedo and Cayot 1994) of geckos in the Galápagos.

SCIENTIFIC NAME	ORIGIN	ACTIVITY	ISLAND(S) OF OCCURRENCE
<i>Gonatodes caudiscutatus</i>	Introduced	Diurnal	San Cristóbal
<i>Hemidactylus frenatus</i>	Introduced	Nocturnal	Isabela
<i>Lepidodactylus lugubris</i>	Introduced	Nocturnal	Isabela, San Cristóbal, Santa Cruz
<i>Phyllodactylus barringtonensis</i>	Endemic	Nocturnal	Islote Santa Fe, Santa Fe
<i>Phyllodactylus bauri</i>	Endemic	Nocturnal	Española, Floreana, Gardner near Española
<i>Phyllodactylus darwini</i>	Endemic	Nocturnal	San Cristóbal
<i>Phyllodactylus galapagoensis</i>	Endemic	Nocturnal	Baltra, Bartolomé, Cowley, Daphne, Fernandina, Isabela, Islote Mares, Pinzón, Plaza Norte, Plaza Sur, Santa Cruz, Santiago, Seymour, Tortuga
<i>Phyllodactylus gilberti</i>	Endemic	Nocturnal	Wolf
<i>Phyllodactylus leei</i>	Endemic	Nocturnal	San Cristóbal
<i>Phyllodactylus reissii</i>	Introduced	Nocturnal	Santa Cruz, Isabela



Hemidactylus frenatus
QCAZ 11128



Lepidodactylus lugubris
QCAZ 10784



Phyllodactylus galapagoensis
QCAZ 11182



Phyllodactylus reissii
QCAZ 10813

FIGURE 1. Species of geckos from Isabela Island, Galápagos. Photographs: O. Torres-Carvajal.

ACKNOWLEDGMENTS: We thank S. Báez, I. Tapia, S. Ron and the 2010 biogeography class of Pontificia Universidad Católica del Ecuador (PUCE) for assistance in the field. We also thank A. Bauer for helpful comments, and the staff of Galapagos National Park in Isabela for help with logistics. This work was funded by PUCE (G19126).

LITERATURE CITED

Bauer, A.M. and K. Henle. 1994. *Das Tierreich 109. Gekkonidae. Part 1, Australia and Oceania*. Berlin: Walter De Gruyter Publishers. 306 p.

Bungartz, F., H. Herrera, P. Jaramillo, N. Tirado, G. Jiménez-Uzategui, D. Ruiz, A. Guézou and F. Ziemmeck. 2011. *Charles Darwin Foundation Galapagos species checklist*. Accessible at <http://www.darwinfoundation.org/datazone/checklists/ecological-groups/introduced-vertebrates/>. Captured on 28 February 2011.

Burt, C.E. 1932. Comments on some lizards from Colombia. *Transactions of the American Microscopical Society* 51: 209-216.

Carrión V., C. Sevilla and W. Tapia. 2008. Management of introduced animals in Galapagos. *Galapagos Research* 65: 46-38.

- Carvajal-Campos, A. and O. Torres-Carvajal. 2010. *Hemidactylus mabouia* Moreau de Jonnés, 1818 and *H. frenatus* Schlegel, 1836 in western Ecuador: new records reveal range extension. *Herpetozoa* 23: 90-91.
- Case, T.J., D.T. Bolger and K. Petren. 1994. Invasions and competitive displacement among house geckos in the tropical Pacific. *Ecology* 75: 464-477.
- Dixon, J.R. and R.B. Huey. 1970. Systematics of the lizards of the gekkonid genus *Phyllodactylus* of mainland South America. *Los Angeles County Museum Contributions in Science* 192: 1-78.
- Garman, S. 1892. The reptiles of the Galapagos islands. *Bulletin of the Essex Institute* 24: 1-15.
- Hoogmoed, M.S. 1989. Introduced geckos in Puerto Ayora, Santa Cruz, with remarks on other areas. *Noticias de Galápagos* 47: 12-16.
- Jadin, R.C., M.A. Altamirano, M.H. Yáñez-Muñoz and E.N. Smith. 2009. First record of the common house gecko (*Hemidactylus frenatus*) in Ecuador. *Applied Herpetology* 6: 193-195.
- Lanza, B. 1973. *On some Phyllodactylus from the Galápagos Islands (Reptilia Gekkonidae)*. Firenze: Museo Zoologico dell'Università di Firenze. 34 p.
- Lubin, Y.D. 1984. Changes in the native fauna of the Galápagos Islands following invasion by the little red fire ant, *Wasmannia auropunctata*. *Biological Journal of the Linnean Society* 21: 229-242.
- Mertens, R. 1963. Die Wiederentdeckung der Geckonengattung *Gonatodes* auf den Galapagos. *Senckenbergiana Biologica* 44: 21-23.
- Olmedo, J. and L. Cayot. 1994. Gecos introducidos en los pueblos de Santa Cruz, San Cristóbal e Isabela. *Noticias de Galápagos* 51-53: 15-20.
- Peck, S.B., J. Heraty, B. Landry and B.J. Sinclair. 1998. Introduced Insect Fauna of an Oceanic Archipelago: The Galápagos Islands, Ecuador. *American Entomologist* 44: 218-237.
- Tapia, W., V. Carrión, M. Patry and H. Snell. 2000. *Estado Actual de los vertebrados introducidos a las islas Galápagos y las estrategias aplicadas por el servicio Parque Nacional Galápagos para su manejo*; p. 77-84 In J. Cevallos and C. Falconi (ed.). Informe Galápagos 2000 -2001. Quito, Ecuador: WWF-Fundación Natura.
- Torres-Carvajal, O. 2001. Lizards of Ecuador: checklist, distribution and systematic references. *Smithsonian Herpetological Information Service* 131: 1-35.
- Van Denburgh, J. 1912. Expedition of the California Academy of Sciences to the Galapagos Islands, 1905-1906. VI. The geckos of the Galapagos Archipelago. *Proceedings of the California Academy of Sciences* 1: 405-430.
- Vanzolini, P.E. 1965. On the *Gonatodes* of the Galapagos Islands (Sauria, Gekkonidae). *Papéis Avulsos do Departamento de Zoologia* 17: 17-19.
- Wright, J.W. 1983. The distribution and status of *Gonatodes collaris* in the Galapagos Archipelago. *Herpetological Review* 14: 32.

RECEIVED: March 2011

LAST REVISED: May 2011

ACCEPTED: May 2011

PUBLISHED ONLINE: July 2011

EDITORIAL RESPONSIBILITY: Pedro L. V. Peloso