

Reptilia, Serpentes, Dipsadidae, *Liophis taeniogaster* Jan, 1863: Distribution extension, new state record and geographic distribution map

Frederico Gustavo Rodrigues França* and Elayne da Silva Bezerra

Universidade Federal da Paraíba, Centro de Ciências Aplicadas e Educação, Departamento de Engenharia e Meio Ambiente. CEP 58297-000. Rio Tinto, PB, Brazil.

* Corresponding author. E-mail: frederico@cca.ufpb.br

ABSTRACT: The current work provides a new state record for the snake *Liophis taeniogaster* in Atlantic Forest of northeastern Brazil.

The genus *Liophis* Wagler, 1830 includes more than 30 recognized species and more than 60 subspecies distributed throughout Central and South America (Dixon 1989; Fernandes *et al.* 2002; Esqueda *et al.* 2005). In Brazil, there are currently 20 *Liophis* species (Bérnils 2010) excluding species of the recently resurrected genus *Lygophis* (Zaher 2009; Vidal *et al.* 2010). This genus is divided into some morphological complex groups as the *L. almadensis* group (Dixon and Thomas 1985), the *L. taeniurus* group (Dixon 2000), the *L. miliaris* group (Dixon 1983a), the Eastern Caribbean group (Dixon 1981), and the *L. cobellus* group (Dixon 1983b).

The *Liophis cobellus* group, as currently recognized, includes four species: *L. breviceps* Cope, 1860, *L. cobellus* (Linnaeus, 1758), *L. frenata* (Werner, 1909), and *L. taeniogaster* Jan, 1863 (Fernandes *et al.* 2002). *Liophis breviceps* and *Liophis cobellus* are distributed only in northern portion of Amazon basin, while *Liophis frenata* occurs in central-southwest Brazil, east Paraguay and north Argentina (Dixon 1989; Giraudo 2001). *Liophis taeniogaster* is the most widely distributed species of this group, presenting apparently two disjoint populations, one in northeastern Brazil ranging from south Bahia to Pernambuco and Maranhão, and another one in Amazon Basin until Lomalinda, Colombia, south to Buena Vista, Bolivia, east to the state of Mato Grosso, Brazil (Fernandes *et al.* 2002).

The *Liophis taeniogaster* differs from the other snakes of the *L. cobellus* group by having eight supralabials and more than 16 maxillary teeth (*L. breviceps* has seven supralabials and less than 16 maxillary teeth), usually less than 33 ventral bands (*L. cobellus* has usually more than 43 ventral bands), and less than 168 ventral scales (*L. frenata* has more than 183 ventrals) (Fernandes *et al.* 2002).

On 08 March 2010, we collected a specimen of *Liophis taeniogaster* (Figure 1) in the city of Rio Tinto, state of Paraíba. The individual was recorded being active on an

off road at 07:00 h (6°48'20" S, 35°04'39" W). Despite the snake was recorded in a disturbed area, the city is close to one of the most important reserves of northeast Atlantic forest, the Reserva Biológica Guaribas. This reserve is a patch with 4.028 ha included in the 2 % of the original forest remaining of northeast Atlantic forest (Silva and Tabarelli 2000), and encloses several fauna and flora species characteristic of the region. The Reserva Biológica Guaribas presents two different vegetation formations (Salgado *et al.* 1981): semi-deciduous rainforest (primary and secondary formation) and savanna.



FIGURE 1. *Liophis taeniogaster* from municipality of Rio Tinto, PB, Brazil (Photo by FGRF).

The snake agrees with the description of *L. taeniogaster* given by Fernandes *et al.* (2002): female with 148 mm snout-vent length, 35 mm caudal length, 2 g weight, supralabials 8/8, infralabials 10/10, dorsal scales in 17-17-15 rows; ventrals 130; cloacal divided; paired subcaudals 46. The specimen was collected (collection permits SISBIO 21799-1) and deposited in Coleção Herpetológica da Universidade Federal da Paraíba (CHUFPB, voucher number RT0005). This is the first record of this species for the state of Paraíba, Brazil (Figure 2).




FIGURE 2. Geographic distribution of *Liophis taeniogaster*. Black circles represent the species distribution according to Fernandes *et al.* (2002). The red star represents the municipality of Rio Tinto, Paraíba, Brazil.

ACKNOWLEDGMENTS: The authors are grateful to Marina Pinheiro Kluppel and ICMbio for collection permits in the Rio Tinto region, Daniel S. Fernandes and Reuber Brandão for valuable comments and suggestions on manuscript, and Daniel S. Fernandes for kindly having sent his article.

LITERATURE CITED

Bérnils, R.S. (org.). 2010. Brazilian reptiles – List of species. Electronic Database Accessible at <http://www.sbherpetologia.org.br/>. Sociedade Brasileira de Herpetologia. Captured on 18 November 2010.

Dixon, J.R. 1981. The neotropical colubrid snake genus *Liophis*: the eastern Caribbean complex. *Copeia* 1981(2): 296-304.

Dixon, J.R. 1983a. Taxonomic status of the South American snakes *Liophis miliaris*, *L. amazonicus*, *L. crysostomus*, *L. mossorensis* and *L. purpurans* (Colubridae: Serpentes). *Copeia* 1983 (3): 791-802.

Dixon, J.R. 1983b. The *Liophis cobella* group of the neotropical colubrid snake genus *Liophis*. *Journal of Herpetology* 17(2): 149-165.

Dixon, J.R. 1989. A key and checklist to the neotropical snake genus *Liophis* with country list and maps. *Smithsonian Herpetological Information Service* 79:1-28.

Dixon, J.R. 2000. Ecuadorian, Peruvian, and Bolivian Snakes of the *Liophis taeniurus* complex with descriptions of two new species. *Copeia* (2):482-490.

Dixon, J.R. and R.A. Thomas. 1985. New Species of South American Water Snake (Genus *Liophis*) from Southeastern Brazil. *Herpetologica* 41(3): 259-262.

Esqueda, L.P., M. Natera, E. La Marca and M. Ilija-Fistar. 2005. Nueva especie de serpiente (Reptilia: Colubridae: *Liophis*) de um bosque tropical relictual em El Estado Barinas, Venezuela. *Herpetotropicos* 2(2): 95-103.

Fernandes, D.S., V.J. Germano, R. Fernandes and F.L. Franco. 2002. Taxonomic status and geographic distribution of the lowland species of the *Liophis cobella* group with comments on the species from the Venezuelan tepuis (Serpentes: Colubridae). *Boletim do Museu Nacional, Nova serie, Zoologia* 481: 1-14.

Giraud, A. 2001. *Serpientes de la Selva Paranaense y del Chaco Húmedo*. Buenos Aires: Literature of Latin America. 328 p.

Salgado, O.A., S.J. Filho and L.M.C. Gonçalves. 1981. As regiões fitoecológicas, sua natureza e seus recursos econômicos. Estudo fitogeográfico; p. 485-544 *In Projeto RADAMBRASIL, Levantamento de Recursos Naturais, vol. 23, Folhas SB 24/25, Jaguaribe / Natal*. Rio de Janeiro: Ministerio das Minas e Energia.

Silva, J.M.C. and M. Tabarelli. 2000. Tree species impoverishment and the future flora of the Atlantic forest of northeast Brazil. *Nature* 404: 72-74.

Vidal, N., M. Dewynter and D.J. Gower. 2010. Dissecting the major American snake radiation: A molecular phylogeny of the Dipsadidae Bonaparte (Serpentes, Caenophidia). *Comptes Rendus Biologies* 333: 48-55.

Zaher, H., F.G. Grazziotin, J.A. Cadle, R.W. Murphy, J.C. Moura-Leite and S.L. Bonatto. 2009. Molecular phylogeny of advanced snakes (Serpentes, Caenophidia) with an emphasis on South American Xenodontines: a revised classification and descriptions of new taxa. *Papéis Avulsos de Zoologia* 49(11): 115-153.

RECEIVED: April 2010
 REVISED: July 2010
 ACCEPTED: October 2010
 PUBLISHED ONLINE: November 2010
 EDITORIAL RESPONSIBILITY: Alejandro R. Giraud