

Distribution extension of *Uracentron flaviceps* (Guichenot, 1855) (Reptilia: Squamata): Second record for the state of Acre, Brazil

Marco Antonio de Freitas^{1*}, Daniella Pereira Fagundes de França² and Diogo Veríssimo³

1 Instituto Chico Mendes de Conservação da Biodiversidade (ICMBio), Rua do INCRA S/N. CEP 69932-000. Brasília, AC, Brazil.

2 Universidade Federal do Acre – UFAC, Campus Universitário. BR 364, Km 04, Distrito Industrial. CEP 69915-900. Rio Branco, AC, Brazil.

3 Durrell Institute of Conservation and Ecology, University of Kent, Canterbury, Kent CT2 7NR, United Kingdom.

* Corresponding author. E-mail: philodryas@hotmail.com

ABSTRACT: A female specimen of *Uracentron flaviceps* (Guichenot, 1855) was found during routine work in the Extractive Reserve Chico Mendes in the municipality of Epitaciolândia, state of Acre, Brazil. This is the second record of the species for the state, extending the known distribution of the species by 570 km.

Uracentron flaviceps (Guichenot, 1855), a lizard species of the family Tropiduridae, is distributed throughout the western part of the Brazilian Amazon, Colombia, Ecuador and Peru (Greene 1977; Vitt and Zani 1996). In Brazil the species has been recorded in the states of Amazonas, Rondônia and for the western extreme of the state of Acre, where a single record exists (Avila-Pires 1995; Avila-Pires *et al.* 2009).

This species is largely restricted to lowland tropical forests, occurring on horizontal limbs high in tropical forest trees (Vitt and Zani 1996). The species has been described to prefer trees with thick irregular bark (Fugler 1986) where it lives and shelters in cavities, in the upper part of the tree trunk (Vitt and Zani 1996). It also uses those cavities for egg-laying, with records of up to 14 eggs in a single nest (Dixon and Soini 1986). *Uracentron flaviceps* is thought to feed mainly on ants, although it occasionally feeds on other insects (Greene 1977; Vitt and Zani 1996).

During a visit to the Extractive Reserve Chico Mendes, in the municipality of Epitaciolândia, state of Acre, a female specimen of *Uracentron flaviceps* (Figure 1) was found lying on a fallen log on the outer edge of a forested area (10°50'17" S, 68°33'28" W). The specimen, with an overall length of 130 mm, was collected (Permit ICMBIO/SISBIO-25261-1.) and deposited in the Herpetological Laboratory of the Universidade Federal do Acre, Campus Floresta, in Cruzeiro do Sul, Acre, Brazil (Voucher UFACF 4001).

In all the herpetological collections reviewed, which included the collection of the Federal University of Acre, Campus Floresta in Cruzeiro do Sul, there was only one previous record of this species for the state of Acre, in the municipality of Cruzeiro do Sul (Avila-Pires 1995). With this new record the species distribution is extended approximately 570 kilometers southeast from Cruzeiro do Sul.

This record is especially valuable given the arboreal habits and cryptic nature of *Uracentron flaviceps*, which

place this species amongst the hardest lizards to study in the Amazon region, and make even the most basic ecological data hard to obtain (Harvey and Gutberlet Jr 1998; Vitt and Zani 1996). Systematic surveys should be undertaken in order to improve our understanding of the distribution of this species and other herpetofauna of the state of Acre, Brazil.



FIGURE 1. *Uracentron flaviceps* recorded at the Reserva Extrativista Chico Mendes, in the municipality of Epitaciolândia, state of Acre, Brazil. Photo by Marco Antonio de Freitas.

ACKNOWLEDGMENTS: The authors wish to thank Paulo Sérgio Bernade for confirming the species identification and for keeping the voucher specimen at UFAC.

LITERATURE CITED

- Avila-Pires, T.C.S. 1995. Lizards of Brazilian Amazonia (Reptilia: Squamata). *Zoologische Verhandlungen* 229: 1-706.
- Avila-Pires, T.C.S., L.J. Vitt, S.S. Sartorius and P.A. Zani. 2009. Squamata (Reptilia) from four sites in southern Amazonia, with a biogeographic analysis of Amazonian lizards. *Boletim do Museu Paraense Emílio Goeldi Ciências Naturais* 4(2): 99-118.
- Dixon, J.R. and P. Soini. 1986. *The reptiles of the upper Amazon Basin, Iquitos Region, Peru*. Milwaukee: Milwaukee Public Museum. 154 p.

- Fugler, C.M. 1986. La Estructura de una comunidad herpetológica en las selvas benianas en la estación de sequia. *Ecología en Bolivia* 8: 1-20.
- Greene, H.W. 1977. Lizards of the genus *Uracentron* (Iguanidae) in east-central Colombia. *Herpetologica* 33(2): 256-260.
- Harvey, M.B. and R.L. Gutberlet Jr, 1998. Lizards of the Genus *Tropidurus* (Iguania: Tropiduridae) from the Serranía de Huanchaca, Bolivia: New Species, Natural History, and a Key to the Genus. *Herpetologica* 54(4): 493-520.
- Vitt, L.J. 1993. Ecology of isolated open formation *Tropidurus* (Reptilia: Tropiduridae) in Amazonian lowland rain forest. *Canadian Journal of Zoology* 71: 2370-2390.
- Vitt, L.J. and P.A. Zani, 1996. Ecology of the elusive tropical Lizard *Tropidurus* [= *Uracentron*] *flaviceps* (Tropiduridae) In Lowland Rain Forest of Ecuador. *Herpetologica* 52(1): 121-132.

RECEIVED: May 2011
REVISED: September 2011
ACCEPTED: October 2011
PUBLISHED ONLINE: December 2011
EDITORIAL RESPONSIBILITY: Raúl Maneyro