

Mollusca, Gastropoda, Ellobioidea, *Carychium minimum* O.F. Müller, 1774: Filling gaps. New population record for the State of New York, Northeastern United States.

Alexander M. Weigand* and Adrienne Jochum

Goethe-University, Institute for Ecology, Evolution and Diversity, Phylogeny and Systematics, Siesmayerstraße 70, 60323. Frankfurt am Main, Germany.

* Corresponding author. E-mail: a.weigand@bio.uni-frankfurt.de

ABSTRACT: The current note reports two new populations of the introduced snail *Carychium minimum* O.F. Müller, 1774 at Ithaca, New York, USA. It confirms the naturalization of this species in Northeastern North America, which was previously known on drift material only.

The Carychiidae (Pulmonata, Ellobioidea) constitute one of only a few gastropod lineages that independently migrated onto land (Vermeij and Dudley 2000; Klussmann-Kolb *et al.* 2008). For the region defining the Northeastern United States and Canada, the taxon *Carychium* O.F. Müller, 1774 comprises at least three native and one introduced species (Figure 1) (Pilsbry 1948; Burch and Van Devender 1980; Hubricht 1985; Nekola and Barthel 2002; Grimm *et al.* 2009). The latter single species, the Herald Thorn *Carychium minimum* O.F. Müller, 1774, was originally endemic to Europe. It demonstrates an amphibious ecology, inhabiting permanent moist environments such as riparian zones, meadows and swamps (Watson and Verdcourt 1953; Morton 1955; Egorov 2007). Beyond its natural occurrence in distribution records, *C. minimum* is a familiar greenhouse species in Europe (Leiss and Reischütz 1996; Horsák *et al.* 2004) where it can be found amidst the roots of plants or on the bottom of flowerpots. Although questionable (Roth 1982), the first record of its occurrence in North America was described by Clapp (1912) who documented a population from Quincy, Massachusetts in close vicinity to a greenhouse. For the most part, only a few and scattered reports in North America (Roth 1982; Forsyth 2004) exist, which are predominantly based on empty shells in drift material, *e.g.* at the Grant River, Ontario, Canada (Forsyth *et al.* 2008).

We collected live *C. minimum* specimens from two localities in Ithaca, New York during early July 2009, which are e-vouchered and stored at the Goethe University Frankfurt am Main, Germany. These two populations were sampled at Beebe Lake on the Cornell Campus (42°26'58"N, 76°28'33"W) and near the Buttermilk Falls State Park off route 96B (42°24'03"N, 76°30'17"W), Tompkins County. Their presence confirms the assumption made by Grimm *et al.* (2009) that this introduced species should be found

living in the Northeastern United States and Canada not only in the proximity of greenhouses and nurseries but also in its natural habitat. These individuals were found between layers of moist leaf litter and sticks at a seep by the lake while the latter population inhabited the riparian zone of a creek. Both juveniles and adult specimens were found. These finds verify that this species has well-established populations at these two locations. Hence, *C. minimum* may very well be more widespread and naturalized in Northeastern North America than previously thought.




FIGURE 1. Native (A-C) and introduced (D) species of *Carychium* O.F. Müller, 1774 in Northeastern United States and Canada. A: *Carychium exiguum* Say, 1882; USA, New York, Grimes Glen. B: *Carychium exile* I. Lea, 1842; USA, New York, Watkins Glen. C: *Carychium nannodes* G.H. Clapp, 1905; USA, Tennessee, Overton County. D: *Carychium minimum* O.F. Müller, 1774; USA, New York, Ithaca, Tompkins County.

ACKNOWLEDGMENTS: We are indebted to Frank Hardie for his tremendous help, patience and perseverance in sorting through piles of leaf litter for these minute snails. Special gratitude goes to Cory Holliday of the

Cave and Karst Incentive Program, Nature Conservancy of Tennessee, Nashville, TN for his speleological prowess and sampling help. For financial support, we wish to thank the Vereinigung von Freunden und Förderern der Johann Wolfgang Goethe-Universität Frankfurt am Main e.V. and the Biodiversity and Climate Research Center (BiK-F), Frankfurt am Main.

LITERATURE CITED

- Burch J.B. and A.S. Van Devender. 1980. Identification of eastern North American land snails: The Prosobranchia, Opisthobranchia and Pulmonata (Actophila). *Walkerana* 1: 33-80.
- Clapp, W.F. 1912. *Carychium minimum* Mull. *The Nautilus* 26: 24.
- Egorov, R. 2007. *Treasure of Russian Shells: Ellobioidei (Cassidulidae, Leucophytiidae, Carychiidae)*. Moscow: Colus v.8, 33 p.
- Forsyth, R.G. 2004. *Land Snails of British Columbia*. Royal BC Museum Handbook. Victoria: Royal British Columbia Museum. 188 p.
- Forsyth, R.G., M.J. Oldham and F.W. Schueler. 2008. Mollusca, Gastropoda, Ellobiidae, *Carychium minimum*, and Ferussaciidae, *Cecilioides acicula*: Distribution extension and first provincial records of two introduced land snails in Ontario, Canada. *Check List* 4(4): 449-452.
- Grimm, F.W., R.G. Forsyth, F.W. Schueler and A. Karstad. 2009. *Identifying Land Snails and Slugs in Canada: Introduced Species and Native Genera*. Ottawa: Canadian Food Inspection Agency. 168 p.
- Horsák M., L. Dvořák and L. Juříčková. 2004. Greenhouse gastropods of the Czech Republic: current stage of research. *Malakologičai Tájékoztató* 22: 141-147.
- Hubricht, L. 1985. *The distributions of the native land mollusks of the Eastern United States*. Fieldiana Zoology New Series 24. Field Museum of Natural History Publication 1359. 191 p.
- Klussmann-Kolb, A., A. Dinapoli, K. Kuhn, B. Streit and C. Albrecht. 2008. From sea to land and beyond - new insights into the evolution of euthyneuran Gastropoda (Mollusca). *BMC Evolutionary Biology* 8: 57.
- Leiss, A. and P.L. Reischütz. 1996. Ein Beitrag zur Kenntnis der Molluskenfauna der Gewächshäuser in Wien und Niederösterreich. *Wissenschaftliche Mitteilungen aus dem Niederösterreichischen Landesmuseum* 9: 173-184.
- Morton, J.E. 1955. The Evolution of the Ellobiidae with a discussion of the origin of the Pulmonata. *Proceedings of the Zoological Society of London* 125: 127-168.
- Nekola, J. and M. Barthel. 2002. Morphometric analysis of *Carychium exile* and *Carychium exiguum* in the Great Lakes region of North America. *Journal of Conchology* 37(5): 515-531.
- Pilsbry, H.A. 1948. *Land Mollusca of North America (north of Mexico)*. *The Academy of Natural Sciences of Philadelphia, Monographs* 3 (2-2): 521-1113.
- Roth, B. 1982. European land mollusks in the San Francisco Bay area, California: *Carychium minimum* Müller and the *Arion hortensis* complex. *The Veliger* 24: 342-344.
- Vermeij, G.J. and R. Dudley. 2000. Why are there so few evolutionary transitions between aquatic and terrestrial ecosystems? *Biological Journal of the Linnean Society* 70: 541-554.
- Watson, H. and B.J. Verdcourt. 1953. The two British species of *Carychium*. *Journal of Conchology* 23: 306-324.

RECEIVED: July 2010

REVISED: September 2010

ACCEPTED: October 2010

PUBLISHED ONLINE: October 2010

EDITORIAL RESPONSIBILITY: Inga Ludmila Veitenheimer Mendes