

NOTES ON GEOGRAPHIC DISTRIBUTION

Reptilia, Squamata, Iguanidae, *Stenocercus aculeatus*: Distribution extension and first record for Ecuador

Omar Torres-Carvajal^{1,2}
Amaranta Carvajal-Campos¹

¹ Pontificia Universidad Católica del Ecuador, Escuela de Biología.
Avenida 12 de Octubre y Roca, Apartado 17-01-2184. Quito, Ecuador. E-mail: omartorcar@gmail.com

² National Museum of Natural History, Smithsonian Institution, Department of Vertebrate Zoology.
Washington, DC 20560, USA

Until recently the name *Stenocercus aculeatus* was applied to two different species described as *Leiocephalus aculeatus* O'Shaughnessy 1879 and *Leiocephalus angulifer* Werner 1901 from eastern Ecuador and northeastern Peru, respectively. These two names were synonymized by Burt and Burt (1931) as *Leiocephalus iridescens aculeatus*, which was subsequently recognized as *Ophryoessoides aculeatus* (Etheridge 1966) and later as *Stenocercus aculeatus* (Frost 1992). Upon resurrection of *Stenocercus angulifer*, which occurs in eastern Ecuador, *S. aculeatus* was restricted to the departments of La Libertad, Loreto, and San Martín at elevations of 723-1311 m in northeastern Peru (Torres-Carvajal 2007).

Herein we report the first record of *Stenocercus aculeatus* for Ecuador based on a donated specimen (QCAZ 8121, male, snout-vent length = 93 mm, tail length = 236 mm), which was collected in the surroundings of Zamora (4°4'12" S, 78°57'36" W, 925 m), province of Zamora Chinchipe (Figure 1). This locality lies approximately 276 km NW from the nearest record (AMNH 57083, Departamento Loreto, Peru) of *Stenocercus aculeatus* reported in the literature (Torres-Carvajal 2007). The record reported in this paper increases to 16 the number of species of *Stenocercus* from Ecuador (Torres-Carvajal 2008).




Figure 1. Distribution of *Stenocercus aculeatus*.

NOTES ON GEOGRAPHIC DISTRIBUTION

Acknowledgements

We thank D. Núñez for donating the specimen herein reported, and E. Arbeláez and D. Salazar for assistance in the field. The specimen was deposited at Museo de Zoología (QCAZ), Pontificia Universidad Católica del Ecuador. The field trip where the specimen was obtained was funded by Smithsonian Institution Restricted Endowment Funds granted to K. de Queiroz and O. Torres-Carvajal.

Literature Cited

- Burt, C. E. and M. D. Burt. 1931. South American lizards in the collection of the American Museum of Natural History and Ecology. *Bulletin of the American Museum of Natural History* 61(7):227-395.
- Etheridge, R. 1966. The systematic relationships of West Indian and South American lizards referred to the iguanid genus *Leiocephalus*. *Copeia* 1966:79-91.
- Frost, D. R. 1992. Phylogenetic analysis and taxonomy of the *Tropidurus* group of lizards (Iguania: Tropiduridae). *American Museum Novitates* 3033:1-68.
- Torres-Carvajal, O. 2007. A taxonomic revision of South American *Stenocercus* (Squamata: Iguania) lizards. *Herpetological Monographs* 21:76-178.
- Torres-Carvajal, O. 2008. Reptiles de Ecuador: lista de especies y distribución. Amphisbaenia y Sauria. Version 1.1. Eltronic Database accesible at <http://www.puce.edu.ec/zoologia/vertebrados/reptiliawebec/reptilesecuador/index.html>. Museo de Zoología, Pontificia Universidad Católica del Ecuador, Quito, Ecuador. Captured on 12 July 2008.

Received July 2008

Accepted September 2009

Published online September 2009