

Physalaemus albifrons Spix, 1824 (Anura: Leiuperidae): Distribution and geographic extension

Cristiane Nikely S. Palmeira^{1*}, Ubiratan Gonçalves² and Viviane N. Paz¹

- 1 Faculdade Integrada Tiradentes – FITS, Setor de Zoologia. Avenida Comendador Gustavo Paiva, 5017, Cruz das Almas. CEP 57038-000. Maceió, AL, Brazil.
2 Universidade Federal de Alagoas, Museu de História Natural, Setor de Zoologia. Avenida Aristeu de Andrade, 452, Farol. CEP 57051-090. Maceió, AL, Brazil.
* Corresponding autor. E-mail: crisnikely@yahoo.com.br

ABSTRACT: We report the first record of *Physalaemus albifrons* in the state of Alagoas, in the municipality of Boca da Mata, Brazil.

The genus *Physalaemus* Fitzinger, 1826 is currently composed of 45 species (Frost 2011), with ample geographic distribution in South America, encompassing all the Brazilian regions, Misiones and Entre Ríos in Argentina, eastern Paraguay, Santa Cruz in Bolivia and likely the plains of southern Venezuela (Costa and Jim 2009). *Physalaemus* species are divided into seven groups: the *P. albifrons* species group including four species, the *P. cuvieri* species group including nine species, the *P. deimaticus* species group including three species, the *P. gracilis* species group including five species, the *P. henselii* species group including three species, the *P. signifer* species group including 13 species (Nascimento *et al.* 2005) and the *P. olfersii* species group including eight species (Cassini *et al.* 2010).

Physalaemus albifrons (Spix, 1824, Figure 1) is a species native to Brazil, found in open areas, from the state of Maranhão to the southern part of the state of Minas Gerais (Andrade and Flora 2004). The species is usually found near water, in waterlogged soil with their body partially submerged or on aquatic macrophytes and is strictly nocturnal, initiating vocalization at dusk (Arzabe *et al.* 2005; Vieira *et al.* 2007).

On 23 May 2011 we collected a male specimen of *P. albifrons* in the municipality of Boca da Mata in the state of Alagoas, Brazil (09°41'19" S, 36°12'07" W; elevation 170 m), in the Cariri da Prensa farm. The individual was found in a temporary pond in a sugar cane plantation. The preserved specimen is registered as MUFAL 8793 in the Museum of Natural History, Universidade Federal de Alagoas – UFAL under collecting license 28997-1 (ICMBIO - Instituto Chico Mendes de Conservação da Biodiversidade).

Although previous reports show a wide occurrence of *P. albifrons* in northeastern Brazil (Table 1; Figure 2), the species had never been recorded from the state of Alagoas. The previous closest records to the new locality are Betânia, Pernambuco (255 km airline to the west, see



FIGURE 1. *Physalaemus albifrons* (MUFAL 8793), collected in the Cariri da Prensa farm, municipality of Boca da Mata, state of Alagoas, Brazil. Photo by Ubiratan Gonçalves.



FIGURE 2. Distribution map of *Physalaemus albifrons*. The new locality is indicated by a red dot.

Borges-Nojosa and Santos 2005) and Aracaju, Sergipe (190 km airline to the south, see Oliveira and Junior 2000). Loebmann and Mai (2008) found the species while

inventorying the anurofauna of three coastal municipalities in the state of Piauí (Parnaíba, Luís Correia and Cajueiro da Praia).

TABLE 1. Localities and references used for the *Physalaemus albifrons* distribution.

MUNICIPALITY	STATE	LATITUDE	LONGITUDE	REFERENCE
Barreirinha	Maranhão	04°19'17" S	44°13'24" W	Nascimento et al. 2005
Brejo Santo	Ceará	07°29'19" S	38°58'58" W	Nascimento et al. 2005; Cruz and Pimenta 2004
Fortaleza	Ceará	03°43'08" S	38°32'33" W	Nascimento et al. 2005
Ibiapaba	Ceará	03°20'05" S	40°42'41" W	Loebmann and Haddad 2010
Aracaju	Sergipe	10°55'00" S	37°04'00" W	Oliveira and Junior 2000
Santa Luzia do Itanhý	Sergipe	11°21'08" S	37°26'44" W	Nascimento et al. 2005
São Cristovão	Sergipe	11°00'48" S	37°13'20" W	Nascimento et al. 2005
Itabaiana	Sergipe	10°41'09" S	37°25'27" W	Carvalho et al. 2005
Barreiras	Bahia	12°08'38" S	44°59'35" W	Nascimento et al. 2005
Maracas	Bahia	13°25'45" S	40°26'15" W	Langone 2001
Bom Jesus da Lapa	Bahia	13°15'13" S	43°24'54" W	Nascimento et al. 2005
Juazeiro	Bahia	09°26'21" S	40°30'17" W	Nascimento et al. 2005
Brotas de macaúbas	Bahia	12°00'24" S	42°37'35" W	Oliveira et al. 2010
Lajeado	Minas Gerais	20°58'46" S	46°57'40" W	Nascimento et al. 2005
Mocambinhos	Minas Gerais	15°51'16" S	43°03'53" W	Langone 2001
Manga	Minas Gerais	14°45'34" S	43°56'02" W	Nascimento et al. 2005
Porteirinha	Minas Gerais	15°44'48" S	43°01'27" W	Nascimento et al. 2005
Serranópolis	Minas Gerais	15°48'53" S	42°51'56" W	Leite et al. 2008
Ilha Grande	Piauí	02°50'05" S	41°49'10" W	Silva et al. 2007
Cabaceiras	Paraíba	07°29'22" S	36°17'18" W	Vieira et al. 2007
São João do cariri	Paraíba	07°23'27" S	36°32'02" W	Vieira et al. 2007
São José do Bonfim	Paraíba	07°09'56" S	37°18'20" W	Vieira et al. 2007
Betânia	Pernambuco	08°16'30" S	38°02'11" W	Borges-Nojosa and Santos 2005
Floresta	Pernambuco	08°38'42" S	38°35'05" W	Borges-Nojosa and Santos 2005

ACKNOWLEDGMENTS: To Professor Célio F. B. Haddad and Carla Cassini for identifying *Physalaemus albifrons*, to Filipe Augusto Nascimento and for valuable suggestions that improved the manuscript, and to Philippe J. R. Kok for additional corrections to the manuscript.

LITERATURE CITED

Andrade, G. and F. Juncá. 2004. *Physalaemus albifrons*. IUCN Red List of Threatened Species. Version 2011.1. Electronic Database accessible at www.iucnredlist.org. Captured on 01 June 2011.

Arzabe, C., G. Skuk, G.G. Santana, F.R. Delfim, Y.C.C. Lima and S.H.F. Abrantes. 2005. Herpetofauna da área de Curimataú, Paraíba; p. 259-274. In F.S. Araújo, M.J.N. Rodal and M.R.V. Barbosa (ed.). *Análise das variações da biodiversidade do Bioma Caatinga: suporte a estratégias regionais de conservação*. Brasília: Ministério do Meio Ambiente.

Borges-Nojosa, D.M. and E.M. Santos. 2005. Herpetofauna da área de Betânia e Floresta, Pernambuco; p. 277-294. In F.S. Araújo, M.J.N. Rodal and M.R.V. Barbosa (ed.). *Análise das variações da biodiversidade do bioma Caatinga: suporte e estratégias de conservação*. Brasília: Ministério do Meio Ambiente.

Carvalho, C.M. and J.C.Vilar. 2005. Introdução - Levantamento da Biota do Parque Nacional Serra de Itabaiana; p.9-14 In C.M. Carvalho and J.C. Vilar (coord.). *Parque Nacional Serra de Itabaiana - Levantamento da Biota*. Aracaju: Ibama, Biologia Geral e Experimental - UFS.

Cassini, C.S., C.A.G. Cruz and U. Caramaschi. 2010. Taxonomic review of *Physalaemus olfersii* (Lichtenstein & Martens, 1856) with revalidation of *Physalaemus lateristriga* (Steindachner, 1864) and description of two new related species (Anura: Leiuperidae). *Zootaxa* 2491: 1-33.

Costa, W.P. and J. Jim. 2009. Composição, diversidade e ocupação espaço-temporal da anurofauna de uma localidade em Anhembi, SP. Electronic Database accessible at http://prope.unesp.br/xxi_cic/27_30769941826.pdf. Captured on 20 June 2011.

Cruz, C.A.G. and B.V.S. Pimenta. 2004. New Species of *Physalaemus* Fitzinger, 1826 From Southern Bahia, Brazil (Anura, Leptodactylidae). *Journal of Herpetology* 38(4): 480-486.

Frost, D.R. 2011. *Amphibian species of the world: an online reference version 5.5 (31 May, 2011)*. Electronic Database accessible at <http://research.amnh.org/herpetology/amphibian/index.php>. Captured on 31 May 2011.

Langone, J.A. 2001. Revisión sobre la distribución geográfica de *Physalaemus albifrons* (Spix, 1824) (Anura: Leptodactylidae). *Cuadernos de Herpetología* 15(1): 85-88.

Leite, F.S.F., F.A. Junca and P.C. Eterovick. 2008. Status do Conhecimento, Endemismo E Conservação de Anfíbios Anuros da Cadeia do Espinhaço, Brasil. *Megadiversidade* 4(1-2): 182-200.

Loebmann, D. and C.F.B. Haddad. 2010. Amphibians and Reptiles from a highly diverse area of the Caatinga Domain: Composition and Conservation Implications. *Biota Neotropica* 10(3): 227-256.

Loebmann, D. and A.C.G. Mai. 2008. Amphibia, Anura, Leiuperidae, *Physalaemus cicada*: Distribution extension in the state of Ceará, Brazil. *Check List* 4(4): 392-394.

Nascimento, L.B., U. Caramaschi and C.A.G. Cruz. 2005. Taxonomic review of the species groups of the Genus *Physalaemus* Fitzinger, 1826 with revalidation of the Genera *Engystomops* Jiménez-De-La-Espada, 1872 and *Eupemphix* Steindachner, 1863 (Amphibia, Anura, Leptodactylidae). *Arquivos do Museu Nacional* 63(2): 297-320.

Oliveira, F.F. and G.P.L. Junior. 2000. Anfíbios Anuros do Campus da Universidade Federal de Sergipe. *Biologia Geral e Experimental* 1(1): 42-74.

Oliveira, M.I.R.R., L.N. Weber and J. Ruggeri. 2010. The tadpole of *Physalaemus albifrons* (Spix, 1824) (Anura, Leiuperidae). *South American Journal of Herpetology* 5(3): 249-254.

Silva, G.E.R., C.L. Santos, M.R. Alves, S.D.V. Sousa and B.B. Annunziata. 2007. Anfíbios das dunas Litorâneas do Extremo Norte do Estado do Piauí, Brasil. *Sitientibus Série Ciências Biológicas* 7(4): 334-340.

Vieira, W.L.S., C. Arzabe and G.G. Santana. 2007. Composição e distribuição espaço-temporal de Anuros no Cariri Paraibano, Nordeste do Brasil. *Oecologia Brasiliensis* 11(3): 383-396.

RECEIVED: August 2011

LAST REVISED: November 2011

ACCEPTED: November 2011

PUBLISHED ONLINE: December 2011

EDITORIAL RESPONSIBILITY: Philippe J. R. Kok