

New distributional records of Gerromorpha Popov, 1971 (Insecta: Hemiptera: Heteroptera) from Cuba

Senén Muñoz Riviaux¹, Carlos Naranjo López² and Felipe Ferraz Figueiredo Moreira^{3*}

- 1 Centro de Aplicaciones Tecnológicas para el Desarrollo Sostenible, Departamento de Gestión Ambiental. Ignacio Agramante 818 e/n Aguilera y Prado, CP: 95100. Guantánamo, Guantánamo, Cuba.
 - 2 Centro Oriental de Ecosistemas y Biodiversidad, Museo de Historia Natural Tomás Romey. Calle Enramada 601, CP: 90100. Santiago de Cuba, Santiago de Cuba, Cuba.
 - 3 Universidade Federal do Rio de Janeiro, Instituto de Biologia, Departamento de Zoologia, Laboratório de Entomologia. Avenida Carlos Chagas Filho, 373, CCS, bloco A, sala 107, Cidade Universitária. CEP 21941-971. Caixa Postal: 68044. Rio de Janeiro, RJ, Brazil.
- * Corresponding author. E-mail: felipento@hotmail.com

ABSTRACT: Distributional records of fourteen species of Gerromorpha, belonging to the families Gerridae, Hebridae, Hydrometridae, Mesoveliidae, and Veliidae, are reported from Cuba. The following species are recorded for the first time from eastern Cuba: *Rheumatobates clanis* Drake and Harris, 1932, *R. meinerti* Schroeder, 1931, *Trepobates pictus* (Herrich-Schäffer, 1848) (Gerridae), *Hydrometra australis* Say, 1832 (Hydrometridae), and *Platyvelia brachialis* (Stål, 1860) (Veliidae).

The infraorder Gerromorpha has approximately 2,100 known species, distributed into eight families and about 160 genera. At least 515 species have been recorded from the Neotropical Region, and more than 100 are yet to be described (Polhemus and Polhemus 2008). According to Naranjo *et al.* (2010), there is a great gap of knowledge concerning the fauna of Gerromorpha from Cuba, but the eastern region of the country is the richest and most diverse portion of the island. Nineteen species of Gerromorpha, belonging to five families and eleven genera, have already been recorded from eastern Cuba, some of which are endemic to the country (Muñoz *et al.* 2010).

The faunistic district of eastern Cuba (de la Cruz 1989) is characterized by its high diversity of landscapes, water bodies dependent of raining seasons, and a high degree of species endemism (Muñoz 1989). The first author of the present study made several collections of aquatic Heteroptera in such area between 2008 and 2010, in both lentic and lotic environments. All material is deposited in the invertebrate collection of the Museo de Historia Natural Tomás Romay, and the records obtained are herein presented. The following abbreviations are used for Cuban faunistic districts and provinces: western Cuba (W) (Isla de la Juventud – IJ, Pinar del Río – PR, Mayabeque – MY); Central Cuba (C) (La Habana – HB, Matanzas – MT, Villa Clara – VC, Sancti Spiritus – SS, Cienfuegos – CF, Camagüey – CM); Eastern Cuba (E) (Holguín – HG, Granma – GR, Santiago de Cuba – SC, Guantánamo – GT). On the lists of examined material, the letter “m” refers to male specimens, the letter “f” to females, and the letter “n” to nymphs.

The distributional data for each species in Cuban territory is presented according to the records listed by Muñoz *et al.* (2010). Distributional expansions are presented for 14 species known from Cuba, and the following species are recorded for the first time from the eastern portion of the country: *Rheumatobates clanis* Drake and Harris, 1932, *R. meinerti* Schroeder, 1931, *Trepobates pictus* (Herrich-Schäffer, 1848) (Gerridae),

Hydrometra australis Say, 1832 (Hydrometridae), and *Platyvelia brachialis* (Stål, 1860) (Veliidae).

Family Gerridae

Limnogonus franciscanus (Stål, 1859)

Known distribution in Cuba. C (HB, CM), E (HG, GR, SC, GT). New records from Cuba. GT – Río Maya, station 2, 20°14'52" N, 74°8'45" W, 15.I.2010: 3 m, 1 f. Río Maya, station 1, 20°14'9" N, 74°9'0" W, 9.VIII.2009: 9 n, 4 m, 11 f. Río Sabanalamar, station 1, 20°11'32" N, 74°48'0" W, 26.IX.2009: 3 n, 2 m, 1 f. Río Sabanalamar, station 2, 20°11'32" N, 74°48'0" W, 14.V.2010: 16 n, 6 m, 6 f. Río Bano, 20°11'18" N, 75°14'9" W, 18.VI.2008: 1 m. Río Macambo, station 2, 20°4'52" N, 74°45'16" W, 17.IV.2010: 7 n, 5 m, 1 f. Río Yacabo, station 1, 20°4'16" N, 74°40'43" W, 24.X.2008: 1 n, 2 m. Laguna Cardonal, station 1, 20°3'49" N, 74°47'15" W, 1.VII.2009: 16 n, 2 m, 2 f. Laguna Cardonal, station 2, 20°3'40" N, 74°46'56" W, 1.VII.2009: 1 m, 1 f. Laguna Macambo, 20°3'36" N, 74°44'46" W, 29.IV.2009: 5 m. Río Macambo, station 1, 20°3'27" N, 74°44'9" W, 25.X.2008: 4 n. Río Imías, station 1, 20°3'22" N, 74°46'32" W, 9.VIII.2010: 11 n, 3 m, 2 f. Río Yacabo, station 2, 20°3'20" N, 74°41'46" W, 4.X.2008: 3 n, 2 m, 1 f. Estuario Playa Imías, 20°3'18" N, 74°36'38" W, 9.VIII.2010: 2 m, 2 f. Estuario Macambo, 20°3'2" N, 74°43'54" W, 20.VI.2009: 3 n, 3 m, 2 f. Río Yateras, station 2, 20°2'17" N, 75°2'10" W, 17.IV.2010: 5 n. Río Yateras, station 1, 19°59'31" N, 74°59'54" W, 8.V.2008: 1 n, 1 m, 1 f. Río Imías, station 2, 18°46'48" N, 74°39'9" W, 1.X.2010: 4 n, 4 m, 3 f.

Metrobates tumidus Anderson, 1932

Known distribution in Cuba. W (MY), C (HB, MT, SS, CM), E (SC, GT).

New records from Cuba. GT – Río Bano, 20°11'18" N, 75°14'9" W, 18.VI.2008: 1 m, 4 f. Río Macambo, station 2, 20°4'52" N, 74°45'16" W, 17.IV.2010: 4 m, 5 f. Río Macambo, station 1, 20°3'27" N, 74°44'9" W, 25.X.2008: 3 m, 2 f. Río Yacabo, station 2, 20°3'20" N, 74°41'46" W, 4.X.2008:

2 m, 7 f. Río Yateras, station 2, 20°2'17" N, 75°2'10" W, 17.IV.2010: 1 n, 2 m, 3 f. Río Yateras, station 1, 19°59'31" N, 74°59'54" W, 8.V.2008: 5 f.

Rheumatobates clanis Drake and Harris, 1932

Known distribution in Cuba. C (MT).

New records from Cuba. GT – Río Maya, station 2, 20°14'52" N, 74°8'45" W, 15.I.2010: 19 n, 2 m, 1 f. Río Yacabo, station 2, 20°3'20" N, 74°41'46" W, 4.X.2008: 3 f. Estuario Hatibonico, 19°53'7" N, 75°19'21" W, 2.X.2009: 18 n, 3 m, 4 f. **[First records from eastern Cuba]**

Rheumatobates meinerti Schroeder, 1931

Known distribution in Cuba. C (CF).

New records from Cuba. GT – Laguna Macambo, 20°3'36" N, 74°44'46" W, 29.IV.2009: 20 n, 13 m, 21 f. Estuario Macambo, 20°3'2" N, 74°43'54" W, 20.VI.2009: 6 n, 2 m, 5 f. **[First records from eastern Cuba]**

Trepobates pictus (Herrich-Schäffer, 1848)

Known distribution in Cuba. C (CF).

New records from Cuba. GT – Arrollo Baitiquirí, 20°2'55" N, 74°53'4" W, 19.IX.2009: 2 m, 1 f. **[First record from eastern Cuba]**

Trepobates taylori (Kirkaldy, 1899)

Known distribution in Cuba. W (IJ), E (GR, HG, SC).

New records from Cuba. GT – Río Macambo, station 2, 20°4'52" N, 74°45'16" W, 17.IV.2010: 3 m, 1 f. Río Imías, station 1, 20°3'22" N, 74°46'32" W, 9.VIII.2010: 1 m, 2 f. Río Yacabo, station 2, 20°3'20" N, 74°41'46" W, 4.X.2008: 6 n, 1 f. **[First records from Guantánamo]**

Family Hebridae

Merragata hebroides White, 1877

Known distribution in Cuba. C (MT), E (SC).

New records from Cuba. GT – Estuario Playa Imías, 20°3'18" N, 74°36'38" W, 9.VIII.2010: 1 f. Río Imías, station 2, 18°46'48" N, 74°39'9" W, 1.X.2010: 1 m, 3 f. **[First records from Guantánamo]**

Family Hydrometridae

Hydrometra australis Say, 1832

Known distribution in Cuba. C (MC).

New records from Cuba. GT – Laguna Cardonal, station 2, 20°3'40" N, 74°46'56" W, 1.VII.2009: 1 m, 1 f. Río Yateras, station 1, 19°59'31" N, 74°59'54" W, 8.V.2008: 2 m. **[First records from eastern Cuba]**

Hydrometra caraiba Guérin-Ménéville, 1857

Known distribution in Cuba. W (PR), C (CF), E (HG).

New records from Cuba. GT – Río Sabanalamar, station 2, 20°11'32" N, 74°48'0" W, 14.V.2010: 2 f. Estuario Macambo, 20°3'2" N, 74°43'54" W, 20.VI.2009: 1 m, 1 f. **[First records from Guantánamo]**

Family Mesoveliidae

Mesovelia mulsanti White, 1879

Known distribution in Cuba. C (HB, MT), E (HG, GR, SC).

New records from Cuba. GT – Río Sabanalamar, station 1, 20°11'32" N, 74°48'0" W, 26.IX.2009: 2 f. Río Yacabo, station 1, 20°4'16" N, 74°40'43" W, 24.X.2008: 4 n, 7 m,

3 f. Laguna Cardonal, station 1, 20°3'49" N, 74°47'15" W, 1.VII.2009: 6 n, 5 m, 1 f. Laguna Cardonal, station 2, 20°3'40" N, 74°46'56" W, 1.VII.2009: 2 m, 8 f. Río Imías, station 1, 20°3'22" N, 74°46'32" W, 9.VIII.2010: 9 n, 5 m, 2 f. Estuario Playa Imías, 20°3'18" N, 74°36'38" W, 9.VIII.2010: 3 f. Arrollo Baitiquirí, 20°2'55" N, 74°53'4" W, 19.IX.2009: 1 m. Río Yateras, station 1, 19°59'31" N, 74°59'54" W, 8.V.2008: 2 n, 2 m. Río Imías, station 2, 18°46'48" N, 74°39'9" W, 1.X.2010: 2 f. **[First records from Guantánamo]**

Family Veliidae

Microvelia cubana Drake, 1951

Known distribution in Cuba. W (IJ), C (HB, CM), E (SC, GT).

New records from Cuba. GT – Río Maya, station 2, 20°14'52" N, 74°8'45" W, 15.I.2010: 4 n, 12 m, 23 f. Río Maya, station 1, 20°14'9" N, 74°9'0" W, 9.VIII.2009: 10 n, 49 m, 83 f. Río Sabanalamar, station 1, 20°11'32" N, 74°48'0" W, 26.IX.2009: 8 n, 15 m, 21 f. Río Macambo, station 2, 20°4'52" N, 74°45'16" W, 17.IV.2010: 29 m, 23 f. Río Yacabo, station 1, 20°4'16" N, 74°40'43" W, 24.X.2008: 9 n, 20 m, 31 f. Laguna Cardonal, station 1, 20°3'49" N, 74°47'15" W, 1.VII.2009: 4 n, 1 f. Laguna Cardonal, station 2, 20°3'40" N, 74°46'56" W, 1.VII.2009: 10 m, 20 f. Río Macambo, station 1, 20°3'27" N, 74°44'9" W, 25.X.2008: 3 n, 3 f. Río Imías, station 1, 20°3'22" N, 74°46'32" W, 9.VIII.2010: 10 n, 35 m, 32 f. Río Yacabo, station 2, 20°3'20" N, 74°41'46" W, 4.X.2008: 3 n, 15 m, 16 f. Estuario Playa Imías, 20°3'18" N, 74°36'38" W, 9.VIII.2010: 8 n, 31 m, 17 f. Estuario Macambo, 20°3'2" N, 74°43'54" W, 20.VI.2009: 5 n, 14 m, 18 f. Arrollo Baitiquirí, 20°2'55" N, 74°53'4" W, 19.IX.2009: 6 n, 25 m, 22 f. Río Yateras, station 2, 20°2'17" N, 75°2'10" W, 17.IV.2010: 4 n, 5 m, 6 f. Río Imías, station 2, 18°46'48" N, 74°39'9" W, 1.X.2010: 15 m, 26 f.

Microvelia longipes Uhler, 1894

Known distribution in Cuba. E (SC).

New records from Cuba. GT – Río Maya, station 1, 20°14'9" N, 74°9'0" W, 9.VIII.2009: 5 m, 20 f. **[First record from Guantánamo]**

Platyvelia brachialis (Stål, 1860)

Known distribution in Cuba. W (PR).

New records from Cuba. GT – Estuario Macambo, 20°3'2" N, 74°43'54" W, 20.VI.2009: 1 f. **[First record from eastern Cuba]**

Rhagovelia collaris (Burmeister, 1835)

Known distribution in Cuba. W (PR), C (HB, VC, SS, CM), E (HG, GR, SC, GT).

New records from Cuba. GT – Río Sabanalamar, station 1, 20°11'32" N, 74°48'0" W, 26.IX.2009: 10 m, 6 f. Río Sabanalamar, station 2, 20°11'32" N, 74°48'0" W, 14.V.2010: 5 m, 12 f. Río Bano, 20°11'18" N, 75°14'9" W, 18.VI.2008: 9 n, 6 m, 4 f. Río Macambo, station 2, 20°4'52" N, 74°45'16" W, 17.IV.2010: 1 m, 2 f. Río Yacabo, station 1, 20°4'16" N, 74°40'43" W, 24.X.2008: 1 n, 4 m, 1 f. Río Macambo, station 1, 20°3'27" N, 74°44'9" W, 25.X.2008: 4 n, 9 m, 5 f. Río Yacabo, station 2, 20°3'20" N, 74°41'46" W, 4.X.2008: 5 n, 10 m, 5 f. Arrollo Baitiquirí, 20°2'55" N, 74°53'4" W, 19.IX.2009: 3 n, 11 m, 14 f. Río Yateras, station 2, 20°2'17" N, 75°2'10" W, 17.IV.2010: 2 m, 6 f. Río Yateras, station 1, 19°59'31" N, 74°59'54" W, 8.V.2008: 5 n, 5 m.

ACKNOWLEDGMENTS: We would like the friend Darwins Rodríguez Mesa for his collaboration during the field work and to the project as a whole, and to José Inacio Ricardo Ribeiro and an anonymous referee for their suggestions to the manuscript. This study benefited from the project CUB/SGP/OP4/CORE/08/04, supported by PPD-GEF.

LITERATURE CITED

- de la Cruz, L. 1989. Regionalización faunística; p. XI.1.2 *In* Nuevo Atlas Nacional de Cuba. Madrid: Instituto de Geografía de la ACC, Instituto Cubano de Geodesia y Cartografía, Instituto Geográfico Nacional de España.
- Muñiz, G.O. 1989. Áreas de alto endemismo y algunas de sus causas; p. X.2.2 *In* Nuevo Atlas Nacional de Cuba. Madrid: Instituto de Geografía de la ACC, Instituto Cubano de Geodesia y Cartografía, Instituto Geográfico Nacional de España.
- Muñoz, S.R., F.F.F. Moreira and C. Naranjo. 2010. Checklist, distribution, and habitat of the semiaquatic and aquatic bugs from Cuba (Hemiptera: Heteroptera: Dipsocoromorpha, Leptopodomorpha, Gerromorpha and Nepomorpha). *Zootaxa* 2562: 1-23.
- Naranjo, C., S.R. Muñoz, F.F.F. Moreira and R.C. Correa. 2010. Aquatic and semiaquatic Heteroptera (Insecta) from Cuba: taxonomic survey and distributional data. *Revista de Biología Tropical* 58: 897-907.
- Polhemus, J.T. and D.A. Polhemus. 2008. Global diversity of true bugs (Heteroptera; Insecta) in freshwater. *Hydrobiologia* 595: 379-391.

RECEIVED: December 2011

ACCEPTED: March 2012

PUBLISHED ONLINE: May 2012

EDITORIAL RESPONSIBILITY: Rodrigo M. Feitosa