

Checklist of benthonic marine invertebrates from Malaga Bay (Isla Palma and Los Negritos), Colombian Pacific

Diego Lozano-Cortés^{1*}, Edgardo Londoño-Cruz^{1,2}, Vanessa Izquierdo¹, Fatnori Arias², Madeleine Barona² and Valentina Zambrano¹

¹ Universidad del Valle, Department of Biology, Coral Reef Ecology Research Group. Calle 13 No. 100 – 00. A.A. 25360, Cali, Colombia.

² Universidad del Valle, Department of Biology, ECOMANGLARES, Estuarine and Mangroves Ecology Research Group, Calle 13 No. 100 – 00. A.A. 25360, Cali, Colombia.

* Corresponding author. Email: diegoffc@gmail.com

ABSTRACT: The composition of marine macroinvertebrates in two localities (Isla Palma and Los Negritos) of Malaga Bay was studied. This bay is located in the Pacific coast of Colombia and was recently declared National Natural Park. The rapid ecological assessment revealed a total of 128 species belonging to 64 families and 11 classes (including threatened species). The most common groups were: Gastropoda (27%), Polychaeta (26%), Malacostraca (16%) and Pelecypoda (13%). Even though the two sites are relatively near, being apart only by 6 km, the composition of the respective communities was very different. They shared only 7.8% of the species found; which might have profound and interesting effects for conservation purposes.

INTRODUCTION

A checklist of regional marine species has many uses. Further to providing base line information and important data for comparative studies on biodiversity it also plays an important role and is an important tool in the recognition and delimitation of areas with need protecting, in the process to infer damage and impacts of anthropogenic activity, in the assessment of the complexity of biological communities, and in the estimation of resource availability (Hendrickx and Harvey 1999).

It is known that comprehensive evaluations and inventories serve as the basis for local practices of conservation, which is relevant in poorly known areas in the tropics (Hendrickx and Harvey 1999). This is the case of Malaga Bay (recently established as a National Natural Park), where the few surveys that have been carried out in the area have shown a rich diversity of intertidal and shallow subtidal marine organisms (Lazarus-Agudelo and Cantera-Kintz 2007). Furthermore, the majority of studies performed on Malaga Bay have been focused on specific taxonomic groups, like mollusks (Blanco and Cantera 1994; Cantera *et al.* 1998), echinoderms (Neira and Cantera 2005; Lozano-Cortés *et al.* 2011), fishes (Castellanos-Galindo *et al.*, 2006), and crustaceans (Lazarus-Agudelo and Cantera-Kintz 2007). In order to have an integral view of the real biodiversity in the zone, the community as a whole should be considered.

Although several areas with high species richness in the world have recently received a lot of attention regarding species censuses, tropical and subtropical marine invertebrate communities are still greatly undescribed. It is important to note that most of the effort has been put on west Indopacific, Tropical Atlantic (TA) and Tropical Eastern Pacific (TEP) (Hendrickx and Harvey 1999). However, the knowledge regarding Colombian marine biodiversity, which has coast in both TA and TEP is scarce and checklists of some groups like worms, bryozoans,

echinoderms and crustaceans are, at best, incomplete. Additionally, it has been suggested that the number of species recorded for these groups represent only 50% of the total richness expected for Colombia (Díaz and Acero 2003).

Considering the incomplete knowledge of the animal biodiversity associated to the intertidal and shallow subtidal marine ecosystems, the aim of this survey was to produce a preliminary checklist of marine macroinvertebrates inhabiting two localities of Malaga Bay (Isla Palma and Los Negritos). Mollusks and crustaceans were sampled as well although they are the most common animals in the intertidal and shallow subtidal habitats. Special attention was put on poorly known taxonomic groups in order to improve the base line knowledge on the biodiversity in the region and to provide useful information to policy makers in order to facilitate decisions regarding the protection and conservation of Malaga Bay.

MATERIALS AND METHODS

Study area

Malaga Bay is located in the middle of the Colombian coast on the Tropical Eastern Pacific (3.93 – 4°08' N; 77.32 – 77°35' W; Figure 1). This area experiences a broad tidal range (*ca.* 4 m) and the weather characteristics are determined by the occurrence of the intertropical convergence zone and the Equatorial low pressure area (Cantera *et al.* 1998; Amaya 2007).

The two sampling localities Isla Palma and Los Negritos, are located at the outer part of the Bay. Isla Palma (3°53' N - 77°21' W; Figure 1) is a small island (*ca.* 138 ha) located less than 4 km off the coast and is surrounded by rocky cliffs with heights between 6 m and 15 m. Los Negritos (3°53' N - 77°24' W; Figure 1) is an intertidal rocky reef located southwest (*ca.* 6 km away) from the coast of Isla Palma and is formed by volcanic rocks in contrast to other rocky areas of Malaga Bay (*e.g.* Isla Palma itself) which are

formed mainly by sedimentary rocks (Lozano-Cortés et al. 2011).

Data collection and identification

At each site, samples were collected by intensive scrutiny of animals hidden in cracks and under stones, both in the subtidal and shallow subtidal, where the sampling was performed using SCUBA and snorkeling. At Isla Palma, the submerged walls of the cliffs were also sampled and some boulders were extracted to look for infauna (e.g. polychaets, mytilid bivalves and alpheid shrimps). At Los Negritos rocky reef, samplings were also conducted during low tide, when the rocks were exposed (intertidal sampling). The maximum depth, at which samplings were performed, was 6 m for Isla Palma and 12 m for Los Negritos.

The samples were kept in seawater until sorting by taxonomic groups was performed. The animals were anesthetized by gradually adding fresh water and magnesium chloride to the seawater. After that, samples were fixed in 10% formalin or 70% alcohol, depending on the group. Photographs of most live or recently fixed animals were taken in order to have a record of natural coloration, since most animals (e.g. nudibranches) lose it when they become in contact with fixatives. Once fixed, samples were transported to the Universidad del Valle marine laboratory for identification and storage. Organisms were identified to the lowest possible taxonomic level following Prahel et al. (1986), Breedy and Guzmán (2003, 2007) for octocorals, Keen (1972) for mollusks, Poupin and Bouchard (2006), Hendrickx (1995), Martin and Davis (2001) and Ng et al. (2008) for crabs, Brusca (1980), Cantera (2011) and FAO guide (Fischer et al. 1995) for other invertebrates. Finally,

all species were preserved in 70% alcohol and stored in the Reference Collection of Marine Biology – Universidad del Valle (CRBMUV for its name in Spanish).

RESULTS AND DISCUSSION

A total of 128 species of marine invertebrates belonging to 64 families were collected (Table 1). The faunal composition was distributed in 10 groups, with 64.1% found in Isla Palma and 40.6% in Los Negritos. Only 7.8% (10 species) was shared between the two places. Within the species found, the presence of 3 species considered threatened on Malaga Bay were relevant (2 cnidarians; *Leptogorgia alba*, *Pacifigorgia symbiotica* and 1 mollusc; *Pinctada mazatlanica*, Castellanos-Galindo et al. 2011). Gastropods and Polychaetes were the groups with the highest richness (34 and 33 species, Figure 2); the latter group was composed of species belonging mostly to endofauna and hence was only collected at Isla Palma (Figure 3), a result that is related to the type of substrate in the locality.

In the class Pelecypoda (16 species), the family Mytilidae was the most representative with 5 species, all of which were collected at Isla Palma. The higher species number in this site is explained in terms of their substrate composition: a considerable number of boulders and some of these rocks, which serve as a habitat for infauna, were removed and sampled. The comparison of the study to previous ones (Guevara-Fletcher 2006, Invemar et al. 2006, Guevara-Fletcher et al. 2011) reveals that there are seven species as new records for Isla Palma (*Barbatia reeveana*, *Trigoniocardia biangulata*, *Papyridea aspersa*, *Lithophaga plumula*, *L. attenuata*, *L. spatiosa*, and *Gregariella coarctata*)




FIGURE 1. Study sites (Isla Palma – IP and Los Negritos – LN) in Malaga Bay, Colombian Pacific Ocean (Modified from Castellanos-Galindo and Giraldo 2008).




FIGURE 2. Total number of species by group for benthic marine invertebrates collected in two localities of Malaga Bay.




FIGURE 3. Species richness of principal marine invertebrates groups on Isla Palma (black bars) and Los Negritos (gray bars).



and six for Los Negritos (*B. reeveana*, *T. biangulata*, *Cardita affinis*, *Periglypta multicostata*, *Gastrochaena ovata* and *Pholadidea tubifera*).

Malacostraca was the third group in number of species with a total of 21, belonging to eight families. The site with the highest diversity was Isla Palma with 76.2 % of the species collected from this class (Figure 3); this diversity could be associated with a higher number of microhabitats that can be seen in this area due to the sedimentary origin of the rocks. However, the number of species collected can be strongly influenced by sampling effort. Almost all of the species reported in this work, have been already reported by other authors working in the bay (Invemar et al. 2006, Lazarus-Agudelo and Cantera-Kintz 2007, Guevara-Fletcher et al. 2011). Nevertheless, there is a new report for Isla Palma (*Neogonodactylus stanschi*, Table 1) expanding its range within the bay.

Echinoderms which are recognized to be a key component in the benthic communities (Hendler et al. 1995) were represented with 11 species divided into 3

classes and 8 families. The most representative class was Ophiuroidea with 6 species. This result contrasts with some areas where this class had the lowest richness (e.g. Malpelo Island; Cohen-Rengifo et al. 2009). Nevertheless, this could be explained because this class is the most diverse group of living echinoderms (Giese et al. 1991) and numerically are the most abundant group in the Colombian Pacific (Neira and Cantera 2005). All the species reported in this work were found in Los Negritos; only 5 of them were found in Isla Palma, possibly because this is a rocky reef where echinoderms have a great ecological significance (Lawrence 1987). Finally, even though the two sample sites are relatively close, the macroinvertebrate composition was quite different, suggesting a habitat differentiation process related to habitat composition and heterogeneity. In this order, this kind of inventories not only helps to improve our knowledge about those species inhabiting these areas but also provides tools for taking decisions to protect those ecosystems that are in need of conservation.

TABLE 1. Checklist of benthic marine invertebrates registered for two localities of Malaga Bay (Isla Palma and Los Negritos). Species marked with an asterisk indicate new reports for the bay (*).

PHYLUM	CLASS	ORDER	FAMILY	SPECIES	LOCALITY				
					Isla Palma	Los Negritos			
PORIFERA	DEMOSPONGIAE	Halichondriida	Hymeniacionidae	<i>Hymeniacion</i> sp. (Bowerbank, 1858)		X			
		Haplosclerida	Haliclonidae	<i>Haliclona</i> sp. (Grant, 1836)		X			
		Leucosdeniida	Grantiidae	<i>Leucandra</i> sp. (Haeckel, 1872)		X			
		Verongiida	Aplysinidae	-		X			
		Hadromerida	Suberitidae	<i>Suberites</i> sp. (Nardo, 1833)		X			
		Axinellida	Axinellidae	-		X			
CNIDARIA	ANTHOZOA	Alcyonacea	Gorgoniidae	<i>Pacificorgia symbiotica</i> (Williams and Breedy, 2004)		X			
				<i>Pacificorgia</i> sp. (Bayer, 1951)		X			
MOLLUSCA	POLYPLACOPHORA	Neoloricata	Ischnochitonidae	<i>Leptogorgia alba</i> (Duchassaing and Michelotti, 1864)	X				
				<i>Radiella dispar</i> (Sowerby, 1832)		X			
	GASTROPODA	Neogastropoda	Acanthochitonidae		<i>Stenoplax limaciformis</i> * (Sowerby, 1832)		X		
					<i>Acanthochitona hirudiniformis</i> (Sowerby, 1832)		X		
			Columbellidae		<i>Columbella strombiformis</i> (Lamarck, 1822)	X			
					<i>Mitrella elegans</i> (Dall, 1871)	X			
			Buccinidae		<i>Cantharus gemmatus</i> (Reeve, 1846)	X	X		
					<i>Cantharus ringens</i> (Reeve, 1846)	X			
			Fascioliariidae		<i>Pustulatirus mediamericus</i> (Hertlein and Strong, 1951)		X		
					<i>Leucozonia cerata</i> (Wood, 1828)	X			
			Muricidae		<i>Murexiella vittata</i> (Broderip, 1836)	X			
					<i>Nucella melones</i> (Duclos, 1832)		X		
					<i>Muricopsis zeteki</i> (Hertlein and Strong, 1951)		X		
					<i>Thais speciosa</i> (Valenciennes, 1832)		X		
					<i>Acanthais brevidentata</i> (Wood, 1828)		X		
					Turbinidae		<i>Turbo saxosus</i> (Wood, 1828)		X
							<i>Trivia pacifica</i> (Sowerby, 1832)	X	
					Cerithiidae		<i>Cerithium uncinatum</i> (Gmelin, 1791)		X
							<i>Cerithium nicaraguense</i> (Pilsbry and Lowe, 1932)		X
					Mitridae		<i>Mitra tristis</i> (Broderip, 1836)		X
<i>Lottia</i> sp. (Gray, 1833)	X								
Lottiidae		<i>Lottia pediculus</i> (Philippi, 1846)				X			
		<i>Nerita scabricosta</i> (Lamarck, 1822)		X					
Cypraeidae		<i>Cypraea robertsi</i> (Hidalgo, 1960)		X					
		<i>Oliva splendidula</i> (Sowerby, 1825)	X						

TABLE 1. CONTINUED.

PHYLUM	CLASS	ORDER	FAMILY	SPECIES	LOCALITY	
					Isla Palma	Los Negritos
MOLLUSCA	GASTROPODA	Neogastropoda	Olivellidae	<i>Olivella rehderi</i> (Olsson, 1956)		X
			Bursidae	<i>Bursa corrugata corrugata</i> (Perry, 1811)		X
				<i>Bursa</i> sp. (Röding, 1798)		X
			Calyptraeidae	<i>Crepidula</i> sp. (Lamarck, 1799)	X	
			Turbinidae	<i>Tegula</i> sp. (Lesson, 1832)	X	
		Sacoglossa	Strombidae	<i>Strombus gracilior</i> (Sowerby, 1825)	X	
			Elysiidae	<i>Elysia diomedea</i> (Bergh, 1894)	X	
		Nudibranchia	Chromodorididae	<i>Chromodoris sedna</i> * (Marcus and Marcus, 1967)		X
				<i>Hypselodoris californiensis</i> * (Bergh, 1879)	X	
				<i>Hypselodoris</i> sp. (Stimpson, 1855)	X	
		Pleurobranchomorpha	Pleurobranchidae	<i>Berthellina ilisima</i> (Marcus and Marcus, 1967)		X
		Anaspidea	Aplysiidae	<i>Dolabrifera dolabrifera</i> (Rang, 1828)		X
		Systellommatophora	Onchidiidae	<i>Onchidella hildae</i> (Hoffmann, 1928)		X
		PELECYPODA	Arcoida	Arcidae	<i>Barbatia reeveana</i> (d'Orbigny, 1846)	X
	<i>Barbatia gradata</i> (Broderip and Sowerby, 1829)				X	X
	Carditoida		Carditidae	<i>Cardita affinis</i> (Sowerby, 1833)		X
	Euheterodonta		Gastrochaenidae	<i>Gastrochaena ovata</i> (Sowerby, 1834)	X	
	Myoida		Pholadidae	<i>Pholadidea tubifera</i> (Sowerby, 1834)	X	
				<i>Lithophaga plumula</i> (Hanley, 1844)	X	X
	Mytiloida		Mytilidae	<i>Lithophaga attenuata</i> (Deshayes, 1836)	X	
				<i>Lithophaga aristata</i> (Dillwyns, 1817)	X	
				<i>Lithophaga spatiosa</i> (Carpenter, 1857)	X	
				<i>Gregariella coarctata</i> (Carpenter, 1857)	X	
		<i>Pinctada mazatlanica</i> (Hanley, 1856)		X		
	Pterioida	Pteriidae	<i>Trigoniocardia biangulata</i> (Broderip and Sowerby, 1829)		X	
			<i>Trachycardium senticosum</i> (Sowerby, 1833)	X		
			<i>Papyridea aspersa</i> (Sowerby, 1833)	X	X	
			<i>Tivela byronensis</i> (Gray, 1838)		X	
	Veneroida	Veneridae	<i>Periglypta multicostata</i> (Sowerby, 1835)		X	
	CEPHALOPODA	Octopoda	Octopodidae	-		X
	ECHINODERMATA	ASTEROIDEA	Valvatida	Ophiasteridae	<i>Pharia pyramidata</i> (Gray, 1840)	X
Ophioplepididae				<i>Ophioplepis plateia</i> (Ziesenhenné, 1940)		X
OPHIUROIDEA		Ophiurida	Ophiocomidae	<i>Ophiocoma aethiops</i> (Lütken, 1859)	X	X
				<i>Ophiocoma alexandri</i> (Lyman, 1860)		X
			Ophiodermatidae	<i>Ophioderma panamensis</i> (Lütken, 1859)		X
				<i>Ophioderma teres</i> (Lyman, 1860)	X	X
ECHINOIDEA		Cidaroida	Cidaridae	<i>Ophiotrix spiculata</i> (Le Conte, 1851)		X
				<i>Hesperocidaris asteriscus</i> (Clark, 1948)	X	X
		Diadematoidea	Diadematidae	<i>Centrostephanus coronatus</i> (Verrill, 1867)	X	X
				<i>Diadema mexicanum</i> (Agassiz, 1863)		X
				<i>Echinometra vanbrunti</i> (Agassiz, 1863)		X
ANNELIDA	POLYCHAETA	Eunicida	Eunicidae	<i>Eunice aphroditois</i> (Pallas, 1788)	X	X
				<i>Eunice</i> sp. 1 (Cuvier, 1817)	X	
				<i>Eunice</i> sp. 2	X	
				<i>Eunice</i> sp. 3	X	
				<i>Eunice</i> sp. 4	X	
				<i>Eunice</i> sp. 5	X	
				<i>Eunice</i> sp. 6	X	
				<i>Eunice</i> sp. 7	X	
				<i>Eunice</i> sp. 8	X	
		<i>Eunice</i> sp. 9	X			
		Canalipalpata	Serpulidae	<i>Serpula</i> sp. (Linnaeus, 1758)	X	
		Amphinomida	Amphinomidae	<i>Amphinomida</i> sp. 1 (Savigny, 1818)	X	
				<i>Amphinomida</i> sp. 2	X	
		Phyllodocida	Syllidae	<i>Syllidae</i> sp. 1 (Grube, 1850)	X	
<i>Syllidae</i> sp. 2	X					
<i>Syllidae</i> sp. 3	X					
		<i>Syllidae</i> sp. 4	X			

TABLE 1. CONTINUED.

PHYLUM	CLASS	ORDER	FAMILY	SPECIES	LOCALITY	
					Isla Palma	Los Negritos
ANNELIDA	POLYCHAETA	Phyllodocida	Nereidae	<i>Platynereis</i> sp. (Kinberg, 1865)	X	
				<i>Nereidae</i> sp. 1 (Fauchald, 1977)	X	
				<i>Nereidae</i> sp. 2	X	
				<i>Nereidae</i> sp. 3	X	
			Phyllodocidae	<i>Nereidae</i> sp. 4	X	
				<i>Eulalia</i> sp. 1 (Savigny, 1818)	X	
				<i>Eulalia</i> sp. 2	X	
				<i>Eulalia</i> sp. 3	X	
			Polynoidae	<i>Eulalia</i> sp. 4	X	
				<i>Polynoidae</i> sp. 1 (Malmgren, 1867)	X	
				<i>Polynoidae</i> sp. 2	X	
				<i>Polynoidae</i> sp. 3	X	
				<i>Polynoidae</i> sp. 4	X	
				<i>Polynoidae</i> sp. 5	X	
			Pilargidae	<i>Polynoidae</i> sp. 6	X	
				<i>Pilargidae</i> sp. (de Saint-Joseph, 1899)	X	
ARTHROPODA	MALACOSTRACA	Decapoda	Alpheidae	<i>Alpheus panamensis</i> (Kingsley, 1878)	X	
				<i>Alpheus</i> sp. (Fabricius, 1798)	X	
			Upogebiidae	<i>Upogebia</i> sp. (Leach, 1814)	X	
			Gonodactylidae	<i>Neogonodactylus stanschi</i> (Schmitt, 1940)	X	
		Diogenidae		<i>Calcinus obscurus</i> (Stimpson, 1859)	X	
			<i>Trizopagurus magnificus</i> (Bouvier, 1898)		X	
		Porcellanidae	<i>Pachycheles calcosus</i> (Haig, 1960)		X	
			<i>Pachycheles chacei</i> (Haig, 1956)	X		
			<i>Pachycheles panamensis</i> (Faxon, 1893)	X		
			<i>Pachycheles vicarius</i> (Nobili, 1901)	X		
			<i>Megalobrachium pacificum</i> (Gore and Abele, 1974)	X		
			<i>Megalobrachium garthi</i> (Haig, 1957)	X		
			<i>Pisidia magdalenensis</i> (Glassell, 1936)	X		
			<i>Petrolisthes donadio</i> (Hiller and Werding 2007)	X		
			Majidae	<i>Mithrax tuberculatus</i> (Stimpson, 1860)	X	
				<i>Mithraculus denticulatus</i> (Bell, 1835)	X	
Xanthidae	<i>Cycloxanthops vittatus</i> (Stimpson, 1860)	X				
	<i>Heteractaea lunata</i> (Milne-Edwards and Lucas, 1844)	X	X			
	<i>Platyactaea dovii</i> (Stimpson, 1871)					
	<i>Paractaea sulcata</i> (Stimpson, 1860)					
Epialtidae	<i>Herbstia tumida</i> (Stimpson, 1871)					
TOTAL		64	128	82	52	

ACKNOWLEDGMENTS: The authors want to express their gratitude to Juan Felipe Lazarus, Jaime Cantera and German Bolívar for helping in the identification of crustaceans, mollusks and polychaetes, respectively. Luz Angela López de Mesa helped in the verification of the Polyplacophora and Pelecypoda species. Raúl Neira, Katherine Torres and Melissa Campuzano were of great help during the field and laboratory phases. Fernando Zapata helped us with figure edition, Jodie Hands review the English version and two reviewers improved this manuscript with their comments. Universidad del Valle supported this study.

LITERATURE CITED

- Amaya, V. 2007. *Echinometra vanbrunti* (Echinometridae) como hospedero de relaciones comensalistas en el Pacífico colombiano. *Acta Biológica Colombiana* 12(1): 57-66.
- Blanco, J.F. and J.R. Cantera. 1994. La familia Conidae (Mollusca: Gastropoda) en el Pacífico colombiano. *Boletín Ecotrópica* 27: 19-39.
- Breedy, O. and H.M. Guzmán. 2003. Octocorals from Costa Rica: The genus *Pacificorgia* (Coelenterata: Octocorallia: Gorgoniidae). *Zootaxa* 281: 1-60.
- Breedy, O. and H.M. Guzmán. 2007. A revision of the genus *Leptogorgia* Milne Edwards & Haime, 1857 (Coelenterata: Octocorallia: Gorgoniidae) in the eastern Pacific. *Zootaxa* 1419: 1-90.
- Brusca, R. 1980. *Common Intertidal Invertebrates of the Gulf of California*. Second edition. Tucson: University of Arizona Press. 513 p.
- Cantera, J.R., R. Neira, and C. Ricaurte. 1998. *Bioerosión en la costa Pacífica colombiana: Un estudio de la biodiversidad, la ecología y el impacto de los animales destructores de acantilados rocosos sobre el hombre*. Primera Edición. Bogotá: Fondo FEN Colombia. 135 p.
- Cantera, J.R. 2011. *Estadios de vida vulnerable de organismos marinos de Bahía Málaga. Guía para su identificación*. Programa Editorial. Cali: Universidad del Valle. 144 p.
- Castellanos-Galindo, G.A., and A. Giraldo. 2008. Food resource use in a tropical eastern Pacific tidepool fish assemblage. *Marine Biology* 153: 1023-1035.
- Castellanos-Galindo, G.A., J. A. Caicedo-Pantoja, L.M. Mejía-Ladino and E. Rubio. 2006. Peces marinos y estuarinos de Bahía Málaga, Valle del Cauca, Pacífico Colombiano. *Biota Colombiana* 7(2): 263-282.
- Castellanos-Galindo, G.A., J.R. Cantera, S. Espinosa and L. M. Mejía-Ladino. 2011. Use of local ecological knowledge, scientist's observations and grey literature to assess marine species at risk in a tropical eastern Pacific estuary. *Aquatic Conservation: Marine and Freshwater Ecosystems* 21: 37-48.
- Cohen-Rengifo, M., S. Bessudo and G. Soler. 2009. Echinoderms, Malpelo Fauna and Flora Sanctuary, Colombian Pacific: New reports and distributional issues. *Check List* 5(3): 702-711.
- Díaz, J.M. and A. Acero. 2003. Marine biodiversity in Colombia: achievements, status of knowledge, and challenges. *Gayana* 67(2): 261-274.

- Fischer, W., F. Krupp, W. Schneider, C. Sommer, K.E. Carpenter and V. H. Niem. 1995. *Guía FAO para la identificación de especies para los fines de la pesca. Pacífico centro-oriental*. Volumen I: Plantas e Invertebrados. Roma: FAO. 646 p.
- Guevara-Fletcher, C. E. 2006. *Macrofauna Bentonica Asociada a los Fondos Sumergidos Blandos y Duros de Bahía Málaga, Pacífico Colombiano*. Tesis de Pregrado. Cali-Colombia: Universidad del Valle, Facultad de Ciencias. 78 p.
- Guevara-Fletcher et al. 2011. Benthic macrofauna associated with submerged bottoms of a tectonic estuary in Tropical Eastern Pacific. *Journal of Marine Biology*, 13 pages.doi:10.1155/2011/193759.
- Giese, C. A., J.S. Pearse and V.B. Pearse. 1991. *Reproduction of Marine Invertebrates – Echinoderms and Lophophorates*. Pacific Grove: The Boxwood Press. 359 p.
- Hendler, G., J.E. Miller, D. L. Pawson and P. M. Kier. 1995. *Sea Stars, Sea Urchins, and Allies: Echinoderms of Florida and the Caribbean*. Washington: Smithsonian Institution Press. 390 p.
- Hendrickx, M.E. 1995. Camarones; p 417-564. In W. Fischer, F. Krupp, W. Schneider, C. Sommer, K. E. Carpenter and V. H. Niem (ed.). *Guía FAO para la identificación de especies para los fines de la pesca, Pacífico centro-oriental*. Vol. I. Plantas e invertebrados. Roma: FAO.
- Hendrickx, M. and A. W. Harvey. 1999. Checklist of anomuran crabs (Crustacea: Decapoda) from the Eastern Tropical Pacific. *Belgian Journal of Zoology* 129(2): 363-389.
- Invemar, Univalle and Inciva. 2006. *Valoración de la biodiversidad marina y costera de Bahía Málaga (Valle del Cauca), como uno de los instrumentos necesarios para que sea considerada un Área Protegida*. Informe Científico Final. Cali: INVEMAR-UNIVALLE-INCIVA. 813 p.
- Keen, M. 1972. *Sea shells of Tropical West America*. Second edition. Stanford: Stanford University Press. 1064 p.
- Lawrence, J. M. 1987. *A functional biology of echinoderms*. Baltimore: The Johns Hopkins University Press. 340 p.
- Lazarus-Agudelo, J. and J.R. Cantera-Kintz. 2007. Crustáceos (Crustacea: Sessilia, Stomatopoda, Isopoda, Amphipoda, Decapoda) de Bahía Málaga, Valle del Cauca (Pacífico Colombiano). *Biota Colombiana* 8(2): 221-239.
- Lozano-Cortés, D.F., E. Londoño-Cruz and F.A. Zapata. 2010. Bioerosión de sustrato rocoso por erizos en Bahía Málaga (Colombia), Pacífico Tropical. *Revista de Ciencias* 15: 9-22.
- Martin, J. W. and G. E. Davis. 2001. *An Updated Classification of the Recent Crustacea*. Los Angeles: Natural History Museum of Los Angeles County. 132 p.
- Neira, R. and J. R. Cantera. 2005. Composición taxonómica y distribución de las asociaciones de equinodermos en los ecosistemas litorales del Pacífico Colombiano. *Revista de Biología Tropical* 53(Suppl. 3): 195-206.
- Ng P.K.L., D. Guinot and P.J.F. Davie. 2008. Systema Brachyurorum: Part I. An annotated checklist of extant brachyuran crabs of the world. *The Raffles Bulletin of Zoology, Supplement* 17: 1-286.
- Poupin J. and J. M. Bouchard. 2006. The eastern Pacific species of the genus *Calcinus* Dana, 1851, with description of a new species from Clipperton Island (Decapoda, Anomura, Diogenidae). *Zoosystema* 28(2): 465-486.
- Prahl, H. von., D. Escobar, and G. Molina. 1986. Octocorales (Octocorallia: Gorgoniidae y Plexauridae) de aguas someras del Pacífico Colombiano. *Revista de Biología Tropical* 34(1):13-33.

RECEIVED: March 2012

ACCEPTED: June 2012

PUBLISHED ONLINE: August 2012

EDITORIAL RESPONSIBILITY: Luis Ernesto Arruda Bezerra