

Magnoliophyta, Arly National Park, Tapoa, Burkina Faso

Oumarou Ouédraogo¹, Marco Schmidt^{2,3,4*}, Adjima Thiombiano¹, Karen Hahn^{3,4}, Sita Guinko¹ and Georg Zizka^{2,3,4}

- 1 Université de Ouagadougou, Laboratoire de Biologie et Ecologie Végétales, UFR/SVT. 03 09 B.P. 848 Ouagadougou 09, Burkina Faso.
 - 2 Senckenberg Research Institute, Department of Botany and molecular Evolution. Senckenberganlage 25, 60325. Frankfurt am Main, Germany
 - 3 J.W. Goethe-University, Institute for Ecology, Evolution & Diversity. Siesmayerstr. 70, 60054. Frankfurt am Main, Germany
 - 4 Biodiversity and Climate Research Institute (BiK-F), Senckenberganlage 25, 60325. Frankfurt am Main, Germany.
- * Corresponding author. E-mail: marco.schmidt@senckenberg.de

ERRATUM

In the page 087 there is a mistake in the title (but not in the legend) of **FIGURE 5** and **FIGURE 6**, which should be read as:

We regret these errors.

FIGURE 5. Chorological spectrum of the flora.

■ = Sudanian; □ = Pantropical; ■ = Sudano-zambesian; ■ = Paletropical;
■ = Afrotropical; ■ = Pluriregional African; ■ = Sudano-Guinean
and Guinean-Congolian; ■ = Afro-American; ■ = Afro-Malagasy;
■ = Cultivated; ■ = Cosmopolitan; □ = Unknown.

FIGURE 6. Life form composition: Number of species per life form.

■ = phanerophyte; □ = therophyte; ■ = chamaephyte; ■ = hemicryptophyte;
■ = geophyte; ■ = hydrophyte; ■ = helophyte.

August 2011.