

Culicidae, province of Misiones, northeastern Argentina

José Martín D'Oría¹, Dardo Andrea Martí^{1*} and Gustavo Carlos Rossi²

1 Laboratorio de Genética Evolutiva, Facultad de Ciencias Exactas, Químicas y Naturales. UNaM. CONICET - Félix de Azara 1552, (3300) Posadas, Misiones, Argentina.

2 Centro de Estudios Parasitológicos y de Vectores (CEPAVE) CCT CONICET, La Plata- UNLP Calle 2 N° 584, (1902) La Plata, Argentina.

* Corresponding author. E-mail: darmarti@yahoo.com.ar

ABSTRACT: We present a preliminary list of the Culicidae species collected in seven localities at the province of Misiones, Argentina, since the mosquito fauna of this region is poorly known. Mosquito sampling was carried out during spring-summer, between August 2006 and January 2007, the months of highest abundance of these insects. Collection of specimens was made with a CDC-like trap (Center for Disease Control and Prevention) in peri-urban zones. Sixty-five species were identified. The geographic distribution of *Anopheles darlingi*, *Uranotaenia davis* and *Howardina fulvithorax* is extended.

INTRODUCTION

The province of Misiones has a surface of 30,000 km² and is located between Paraguay and Brazil; it comprises part of the phytogeographic region of the semi-deciduous Atlantic forest in its northern portion and the southernmost area of the Distrito de los Campos. Mean temperatures range from 15 °C in winter to 26 °C in summer, with a mean annual rainfall of 1,800 mm. The remainder of the interior Atlantic forest was recently estimated as 41-55 % of the original area (between 11,000 km² to 17,000 km²). This environment is a moderator of the climate, preserving a high level of humidity in the atmosphere and influencing high biodiversity in the flora, thus favouring the existence of a great diversity of animal species (Giraudo *et al.* 2003).

With the exception of the city of Posadas (Figure 1), all other collection sites in this study were within the area bound on the east by the Parque Estadual do Turvo, state of Rio Grande do Sul, Brazil, and in the north by the Parque Estadual Foz do Iguazu, state of Paraná, Brazil, a region with the lowest anthropic degradation within the geographic area considered (Giraudo *et al.* 2003).

At present, about 3,622 species of mosquitoes are known in the World (Harbach and Kitching 1998; W.R.B.U. 2008). Until now, 226 (6.23 %) of them have been recorded in Argentina, of which 165 were found in the province of Misiones (Rossi *et al.* 2006). Many of

these species are actual or potential vectors of infectious agents that can cause important human diseases (Consoli and Lourenço-de-Oliveira 1994; Forattini 2002). The goal of this study was to update the record of mosquito species in seven localities of the province of Misiones, with emphasis in the eastern region along the Uruguay River coast, where Culicidae fauna was poorly known.

MATERIAL AND METHODS

The sampling localities were near zones of cross-border step, with high harbour and commercial activity and with an international bridge crossing of people between Argentina, Paraguay and Brazil (Figure 1). The sampling localities were (1) Puerto Iguazú (25°35'28" S, 54°33'38" W); (2) Comandante Andresito, with two sampling sites, (2a) Prefectura Naval Argentina post (25°35'25" S, 53°59'20" W), and (2b) Gendarmería Nacional Argentina post (25°36'02" S, 53°58'19" W); (3) Bernardo de Irigoyen (26°16'13" S, 53°39'28" W); (4) El Soberbio (27°17'08" S, 54°11'48" W); (5) Alba Posse with two sampling sites, (5a) Aduana de Alba Posse (27°34'17" S, 54°40'40" W) and (5b) Alba Posse (27°34'10" S, 54°40'43" W); (6) San Javier (27°53'34" S, 55°08'03" W) and (7) Posadas (27°25'11" S, 55°51'41" W).

Capture of adult mosquitoes was carried out with a CDC-like trap, with minor modifications (Sudia and Chamberlain 1962). Samples were obtained monthly from August 2006 to January 2007; the climatic data of the seven sampling localities recorded during the present work are shown in Table 1. Traps were operated during 20 continuous hours in each site. They were set at 1.7 m above ground. The attractive element in this type of trap is carbon dioxide (CO₂-baited), supplied by a cylinder with a gas-flowmeter, calibrated to released 0.4 l/h, and four LED light bulbs. Traps were recovered after the scheduled time, and specimens collected and mounted for identification using taxonomic keys (Darsie 1985).

The abbreviation of genera and subgenera is based on Reinert (2001) *Howardina*, *Ochlerotatus* and *Stegomyia* are considered as genera according to Reinert (2000), as well as *Lutzia* accordingly to Tanaka (2003). Voucher specimens of each species were deposited in the collection of the




FIGURE 1. Location of the study areas. See details in the text.

TABLE 1. Climatic data for the sampling localities of the province of Misiones, Argentina. Mean monthly rainfall (mm) / mean monthly temperature (°C). ND= No data available.

Localities	2006					2007
	August	September	October	November	December	January
(1) Puerto Iguazú	63.0 / ND	47.8 / ND	55.0 / ND	25.6 / ND	26.2 / ND	26.2 / ND
(2) Cte. Andresito	17.4 / 18.7	34.6 / 20.4	16.5 / 25.7	15.6 / 25.4	18.8 / 27.4	24.4 / 27.2
(3) Bernardo de Irigoyen	29.1 / 15.8	17.2 / 18.7	26.7 / 21.5	16.8 / 19.8	26.4 / 23.2	31.5 / 21.8
(4) El Soberbio	21.7 / 16.3	14.2 / 18.7	17.9 / 21.6	32.0 / 21.6	17.6 / 24.7	26.8 / 24.6
(5) Alba Posse	18.2 / ND	13.5 / ND	35.6 / ND	23.8 / ND	25.6 / ND	15.0 / ND
(6) San Javier	28.7 / 15.9	21.1 / 17.6	30.5 / 24.3	24.9 / 24.0	27.1 / 28.3	22.9 / 28.4
(7) Posadas	23.0 / 18.3	20.8 / 18.4	27.1 / 23.9	35.4 / 23.6	19.7 / 28.3	28.8 / 27.7

Museo de La Plata, Argentina (MLP). Authorization for collecting specimens in the protected areas was provided by Dirección de Flora y Fauna, Ministerio de Ecología de Misiones, Argentina.

RESULTS AND DISCUSSION

More than 2,000 specimens were collected, 1,702 were identified (1,656 females and 46 males), corresponding to 65 known species within 18 genera (Table 2); the remaining specimens could not be identified because of the damage produced by the trap. The most diverse tribe was Aedini (21 species) followed by Sabethini (17 species) and Culicini (15 species).

Uranotaenia davisii was collected in December 2006 and January 2007 in the city of El Soberbio. This species was previously reported from the province of Formosa, Argentina, by Duret (1950). *Howardina fulvithorax* was also collected in Eldorado. This is the first report for both species in Misiones.

This preliminary study on the richness of mosquito species carried out during six months, mainly in eastern localities of Misiones, contributes to the knowledge of the mosquito fauna in densely inhabited peri-urban zones. Several species collected during this work are indicated as vectors of infectious agents in the Neotropical region (Table 2); however, there are not enough records in Argentina about the possibility that all these mosquitoes were vectors in the country.

Anopheles albitarsis s.l. is considered a complex of species (Albitarsis Complex), being collected in the sampling area with *An. deaneorum* (Rosa Freitas 1989). Both species can only be reliably identified by means of the clipeal setae in the head of the fourth-instar larva. The other species of the complex have not been mentioned in the area and their identification can only be done through other technical methods, such as molecular DNA analysis (Wilkerson et al. 1995). *Anopheles albitarsis* is considered a secondary vector of malaria in Argentina, while *An. darlingi* is the principal vector of the disease, at least in the north-eastern area of the Country (Bejarano 1960).

Anopheles darlingi is considered the most efficient vector of malaria in the Amazon basin and part of South America (Rubio-Palis and Zimmerman 1997; Rossi and Almirón 2004). The previous record of this species in the province of Misiones was from the Corpus locality (27°7'04" S, 55°32'34" W), near the Paraná River from a single badly preserved mosquito (Tricio et al. 2002). In the present

TABLE 2. Species of Culicidae (Diptera) collected in seven localities of the province of Misiones, Argentina, from August 2006 to January 2007.

Loc = Locality referenced in Figure 1; ♀/♂ = number of females and males specimens collected; * = species indicated as vectors of infectious agents in the Neotropical region.

Taxon	Loc	♀	♂
Family Culicidae			
Subfamily Anophelinae			
Genus <i>Anopheles</i> Meigen			
<i>An. (Anopheles) fluminensis</i> Root	5, 2	11	0
<i>An. (Ano.) mediopunctatus</i> (Theobald)	6	13	0
<i>An. (Nyssorhynchus) albitarsis</i> s.l.* Lynch Arribalzaga	4	4	0
<i>An. (Nys.) darlingi</i> * Root	3	4	0
<i>An. (Nys.) evansae</i> (Brethes)	1, 2, 5	31	0
<i>An. (Nys.) rondoni</i> (Neiva and Pinto)	1, 7, 3, 4	74	2
<i>An. (Nys.) triannulatus</i> (Neiva and Pinto)	3	18	1
Genus Chagasia (Cruz)			
<i>Ch. fajardi</i> (Lutz)	2, 3	40	0
Subfamily Culicinae			
Tribe Aedeomyiini			
Genus Aedeomyia Theobald			
<i>Ad. (Aedeomyia) squamipennis</i> (Lynch Arribalzaga)	1	11	0
Tribe Aedini			
Genus Howardina Theobald			
<i>Howardina fulvithorax</i> (Lutz)	4	2	0
Genus Ochlerotatus Lynch Arribalzaga			
<i>Oc. (Ochlerotatus) albifasciatus</i> * (Macquart)	1	25	0
<i>Oc. (Och.) crinifer</i> * (Theobald)	2	7	0
<i>Oc. (Och.) fluviatilis</i> (Lutz)	4, 5, 7	37	3
<i>Oc. (Och.) fulvus</i> * (Wiedemann)	1	3	1
<i>Oc. (Och.) hastatus</i> Dyar	5, 2, 3, 4	53	1
<i>Oc. (Och.) scapularis</i> * (Rondani)	1, 2, 3	49	2
<i>Oc. (Och.) serratus</i> * s.l. (Theobald)	1, 2, 4-6	67	0
<i>Oc. (Och.) synchytyus</i> Arnell	2	4	0
<i>Och. (Protomacleaya) terreus</i> (Walker)	2-3	8	1
Genus Stegomyia Linnaeus			
<i>St. aegypti</i> * (Linnaeus)	1, 4, 7	6	0
<i>St. albopicta</i> * (Skuse)	1	3	0
Genus Psorophora Robineau-Desvoidy			
<i>Ps. (Janthinosoma) albigena</i> (Peryassu)	4	52	0

TABLE 2. (CONTINUED)

Taxon	Loc	♀	♂
<i>Ps. (Jan.) albipes</i> (Theobald)	2-6	96	5
<i>Ps. (Jan.) cyanescens</i> (Coquillett)	4	4	0
<i>Ps. (Jan.) discrucians</i> (Walker)	2,4	15	0
<i>Ps. (Jan.) ferox*</i> (Von Humboldt)	2-5	116	11
<i>Ps. (Jan.) lutzii</i> (Theobald)	5	5	0
<i>Ps. (Psorophora) ciliata</i> (Fabricius)	4	31	0
<i>Ps. (Pso.) cilipes</i> (Fabricius)	4	27	0
Genus Haemagogus Willinston			
<i>Hg. (Conopostegus) leucocelaenus*</i> (Dyar and Shannon)	1-6	66	0
Tribe Culicini			
Genus Culex Linnaeus			
<i>Cx. (Culex) bidens</i> Dyar	3	5	3
<i>Cx. (Cux.) brethesi</i> Dyar	3	5	0
<i>Cx. (Cux.) chidesteri</i> Dyar	3	7	1
<i>Cx. (Cux.) coronator</i> Dyar and Knab	3-5	51	3
<i>Cx. (Cux.) dolosus</i> Complex Lynch Arribalzaga	7	5	0
<i>Cx. (Cux.) dolosus</i> Lynch Arribalzaga	1-5,7	31	4
<i>Cx. (Cux.) quinquefasciatus*</i> Say	1,3-5,7	27	1
<i>Cx. (Melanoconion) pilosus</i> (Dyar and Knab)	1-3-7	17	4
Genus Lutzia Theobald			
<i>Lutzia bigoti</i> Bellardi	2	7	0
Tribe Mansoniini			
Genus Coquillettidia Dyar			
<i>Cq. (Rhynchotaenia) fasciolata</i> (Lynch Arribalzaga)	3,4	236	3
<i>Cq. (Rhy.) hermanoi</i> (Lane and Coutinho)	4	5	0
<i>Cq. (Rhy.) juxtamanonia</i> (Chagas)	2	4	0
<i>Cq. (Rhy.) nigricans</i> (Coquillett)	3,4	9	0
<i>Cq. (Rhy.) shannoni</i> (Lane and Antunes)	7	3	0
<i>Cq. (Rhy.) venezuelensis</i> (Theobald)	3,4	15	0
Genus Mansonia Blanchard			
<i>Ma. (Mansonia) humeralis</i> Dyar and Knab	7,1	75	0
<i>Ma. (Man.) titillans*</i> (Walker)	7,1	38	0
Tribe Sabethini			
Genus Limatus Theobald			
<i>Limatus durhamii</i> Theobald	1,6,2,4	17	0
Genus Onirion Harbach and Peyton			
<i>On. Brucei</i> (Del Ponte and Cerqueira)	3	6	0
Genus Sabethes Robineau-Desvoidy			
<i>Sa. (Peytonulus) identicus</i> Dyar and Knab	3	9	0
<i>Sa. (Sabethes) albiprivus</i> Theobald	5,6	11	0
<i>Sa. (Sab.) belisarioi</i> Neiva	2	6	0
<i>Sa. (Sab.) purpureus</i> (Theobald)	6	4	0
Genus Trichoprosopon Theobald			
<i>Tr. compressum</i> Lutz	3	22	0
<i>Tr. pallidiventer</i> (Lutz)	3,4,5	35	0
<i>Tr. simile</i> Lane and Cerqueira	3,6	28	0
Genus Wyeomyia Theobald			
<i>Wy. (Phoniomyia) flabellata</i> Lane and Cerqueira	2	9	0
<i>Wy. (Pho.) pilicauda</i> Root	4	5	0
<i>Wy. (Wyeomyia) codiocampa</i> Dyar and Knab	1	7	0

TABLE 2. (CONTINUED)

Taxon	Loc	♀	♂
<i>Wy. (Wyo.) limai</i> Lane and Cerqueira	3	13	0
<i>Wy. (Wyo.) oblita</i> (Lutz)	4,5	7	0
<i>Wy. aporonoma</i> Dyar and Knab	1	16	0
<i>Wy. mystes</i> Dyar	5	9	0
<i>Wy. serratoria</i> (Dyar and Nunez Tovar)	1	21	0
<i>Wyeomyia</i> spp.	1	3	0
Tribe Uranotaeniini			
Genus Uranotaenia Theobald			
<i>Ur. (Uranotaenia) davisii</i> Lane	4	6	0

work, the distribution of *An. darlingi* is extended to Bernardo de Irigoyen (Figure 1), approximately 200 km to the northeast of Corpus city.

Ochlerotatus albifasciatus was incriminated as vector of the Western Equine Encephalitis virus (WEE) in central Argentina, reaching to northern Patagonia (Aviles *et al.* 1992). *Stegomyia aegypti* is vector of Dengue viruses in north-western and northeastern Argentina (Avilés *et al.* 1999). The first report of West Nile virus (WNV) in birds in Argentina was reported in 2004 (Díaz *et al.* 2008), and its isolation from equines in 2006 (Morales *et al.* 2006), *Culex pipiens quinquefasciatus* possibly being the vector in our Country as in other parts of the World.

Another *Flavivirus*, St. Louis Encephalitis virus (SLE) is transmitted by *Cx. pipiens quinquefasciatus* in Argentina (Mitchell *et al.* 1980; Contigiani and Spinsanti 2002; Diaz *et al.* 2003). According to Bejarano (1959a; b), *Haemagogus leucocelaenus* could be vector of Yellow Fever virus (YF). Recently, Vasconcelos *et al.* (2003) reported the detection of YF antigen by immunohistochemistry in the liver samples of dead monkeys from the municipalities of Santo Antonio das Garruchos and Missfies, state of Rio Grande do Sul, Brazil. Unfortunately, there are no data in Argentina on mosquitoes that were responsible for the human cases of Yellow Fever in the rural areas of the municipalities of San Vicente and Eldorado in province of Misiones, reported during April 2008.

We would emphasize the presence of the principal vectors of Yellow Fever or Dengue viruses, *i.e.*, *Haemagogus leucocelaenus*, *Stegomyia aegypti* and *Stegomyia albopicta* in the province. The last species can be found in wild, rural, urban and peri-urban environments and could connect the selvatic and urban cycles of the Yellow Fever (Consoli and Lourenço-de-Oliveira 1994). In this framework, the presence of these vectors deserves major attention by sanitary authorities.

ACKNOWLEDGMENTS: The authors are indebted to Mr. Ernesto R. Krauczuk (Flora and Fauna director MERNRYT) for allowing collection of mosquitoes in protected areas. The research of J.M.D. was supported by a scholarship of CEDyT. The continuous support of Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET) and Universidad Nacional de Misiones (UNaM) is acknowledged. The authors are also grateful to Dr. Claudio J. Bidau for kindly reviewing the manuscript, to Eric Stolar (INTA, Cerro Azul) for climatic data assistance and to Mr. Henry R. Rupp by the English language revision.

LITERATURE CITED

Avilés, G., M. S. Sabattini and C. J. Mitchell. 1992. Transmission of

- Western equine encephalomyelitis virus by Argentine *Aedes (Ochlerotatus) albifasciatus* (Diptera: Culicidae). *Journal of Medical Entomology* (29): 850-853.
- Avilés, G., G. Rangeon, V. Vorndam, A. Briones, P. Baroni, D. Enria and M. S. Sabattini. 1999. Dengue reemergence in Argentina. *Emerging Infectious Diseases* (5): 575-578.
- Bejarano, J. F. R. 1959a. Anophelini de la República Argentina y su relación con el Paludismo. *II Jornadas Entomopidemiológicas Argentinas* (2): 305-329.
- Bejarano, J. F. R. 1959b. Complejo patogénico de la fiebre amarilla en América. *Primeras Jornadas Entomopidemiológicas Argentinas* (1): 653-702.
- Bejarano, J. F. R. 1960. Distribución geográfica del *A. (N.) darlingi* en la República Argentina. *Actas 1º Congreso Sudamericano de Zoología*. Tomo (1): 189-197.
- Consoli, A. G. B. R. and R. Lourenço-de-Oliveira. 1994. *Principais mosquitos de importância sanitária do Brasil*. Rio de Janeiro: Editora FIOCRUZ. 225 p.
- Contigiani, M. and L. Spinsanti. 202. Arbovirus en la Argentina. *Revista de la Sociedad Entomológica Argentina* (Special Publication N°2): 19-22.
- Darsie, R. F. 1985. Mosquitoes of Argentina. Part I. Keys for identification of adult females and fourth stage larvae in English and Spanish (Diptera, Culicidae). *Mosquito Systematics* 17 (3): 153-253.
- Díaz, L. A., W. R. Almirón, F. Ludueña Almeida, L. I. Spinsanti and M. S. Contigiani. 2003. Vigilancia del virus encefalitis de San Luis y mosquitos (Diptera: Culicidae) en la provincia de Córdoba, Argentina. *Entomología y Vectores* 10: 551-566.
- Díaz, L. A., N. Komar, A. Visintin, M. J. Dantur Juri, M. Stein, R. Lobo Allende, L. Spinsanti, B. Königheim, J. Aguilar, M. Laurito, W. Almirón, and M. Contigiani. 2008. West Nile Virus in Birds, Argentina. *Emerging Infectious Diseases* 14: 689-690.
- Duret, J. P. 1950. Contribución al conocimiento de la distribución geográfica de los culicidos argentinos. (Diptera-Culicidae). *Revista de Sanidad Militar Argentina* (49): 365-380.
- Forattini, O. P. 2002. *Culicidologia Médica: identificação, biologia, epidemiologia*. Vol. 2. São Paulo: Editora da Universidade de São Paulo. 864 p.
- Giraud, A. R., H. Povedano, M. J. Belgrano, E. Krauczuk, U. Pardiñas, A. Miquelarena, D. Ligier, D. Baldo, and M. Castelino. 2003. Biodiversity status of the Interior Atlantic Forest of Argentina; p. 160-180 In C. Galindo-Leal and I. G. Cámara (ed.), *The Atlantic Forest of South America*. New York: Island Press.
- Harbach, R. E. and I. J. Kitching. 1998. Phylogeny and classification of the Culicidae (Diptera). *Systematic Entomology* 23: 327-370.
- Mitchell, C. J., T. P. Monath and M. S. Sabattini. 1980. Transmission of St. Louis Encephalitis virus from Argentina by mosquitoes of the *Culex pipiens* (Diptera: Culicidae) complex. *Journal of Medical Entomology* 17: 282-285.
- Morales, M. A., M. Barrandeguy, C. Fabbri, J. B. Garcia, A. Vissani, K. Trono, G. Gutierrez, S. Pigretti, H. M., N. Garrido, N. Taylor, F. Fernandez, S. Levis, and D. Enría. 2006. West Nile Virus Isolation from Equines in Argentina, 2006. *Emerging Infectious Diseases* 12: 1559-1561.
- Reinert, J. F. 2000. New classification for the composite genus *Aedes* (Diptera: Culicidae: Aedini), elevation of subgenus *Ochlerotatus* to generic rank, reclassification of the other subgenera, and notes on certain subgenera and species. *Journal of the American Mosquito Control Association* 16: 175-188.
- Reinert, J. F. 2001. Revised list of abbreviations for genera and subgenera of Culicidae (Diptera) and notes on generic and subgeneric changes. *Journal of the American Mosquito Control Association* 17 (1): 51-55.
- Rosa Freitas, M. G. 1989. *Anopheles (Nyssorhynchus) deaneorum*: A new species in the *albitarsis* Complex (Diptera: Culicidae). *Memórias do Instituto Oswaldo Cruz* 84: 535-543.
- Rossi, G. C. and W. R. Almirón. 2004. Clave ilustrada para la identificación de larvas de mosquitos de interés sanitario encontradas en criaderos artificiales en la Argentina. *Serie: Enfermedades Transmisibles*. Monografía N° 5. Buenos Aires: Ed. Mundo Sano. 53 p.
- Rossi, G. C., E. A. Lestani and J. M. D'Oria. 2006. Nuevos registros y distribución de mosquitos de la Argentina. *Revista de la Sociedad Entomológica Argentina* 65 (3-4): 51-56.
- Rubio-Palis, Y. and R. H. Zimmerman. 1997. Ecoregional classification of malaria vectors in the neotropics. *Journal of Medical Entomology* (34): 499-510.
- Sudia, W. D. and R.W. Chamberlain. 1962. Battery operated light trap, an improved model. *Mosquito News* (22): 126-129.
- Tanaka, K. 2003. Studies on the pupal Mosquitoes of Japan (9) Genus *Lutzia*, with establishment of two new subgenera, *Metalutzia* and *Insulalutzia* (Diptera: Culicidae). *Japanese Journal of Systematic Entomology* (9): 159-169.
- Tricio, A., P. M. Morawicki, C. I. Fernández Díaz, F. Krsticevic, and S. Araki. 2002. Monitoreo de dípteros vectores hematófagos en el área de influencia de la represa de Yacyretá. Período febrero 2000-enero 2001; p 173-183 In O. D. Salomón (ed.), *Actualizaciones en Artropodología Sanitaria Argentina*. Serie *Enfermedades Transmisibles*. Buenos Aires: Fundación Mundo Sano.
- Vasconcelos, P. F. C., A. F. Sperb, H. A. O. Monteiro, M. A. N. Torres, M. R. S. Sousa, H. B. Vasconcelos, L. B. L. F. Mardini and S.G. Rodrigues. 2003. Isolations of yellow fever virus from *Haemagogus leucocelaenus* in Rio Grande do Sul State, Brazil. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 97: 60-62.
- Wilkerson, R. C., T. J. Parsons, T. A. Klein, T. V. Gaffigan, E. Bergo and J. Consolim. 1995. Diagnosis by random amplified polymorphic DNA polymerase chain reaction of four cryptic species related to *Anopheles (Nyssorhynchus) albitarsis* (Diptera: Culicidae) from Paraguay, Argentina and Brazil. *Journal of Medical Entomology* (32): 697-704.
- WRBU, 2008. Walter Reed Biosystematics Unit. Systematics Catalog of Culicidae. Electronic database accessible at <http://www.mosquitocatalog.org/main.asp>. Captured on April 10, 2008.

RECEIVED: July 2009

REVISED: October 2009

ACCEPTED: October 2009

PUBLISHED ONLINE: March 2010

EDITORIAL RESPONSIBILITY: Ana Lúcia Tourinho